

FORM PTO-1390
(REV 5-93)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371

2055MC/48896

U.S. APPLICATION NO. (if known, see 37 CFR 1.5)

09/554414

INTERNATIONAL APPLICATION NO.
PCT/CA98/01059INTERNATIONAL FILING DATE
12 November 1998PRIORITY DATE CLAIMED
12 November 1997TITLE OF INVENTION
DNA DEMETHYLASE, THERAPEUTIC AND DIAGNOSTIC USES THEREOFAPPLICANT(S) FOR DO/EO/US
Mosthe SZYF, Sanjoy BHA TACHARYA; Shyam RAMCHANDANI

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2)).
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ has been transmitted by the International Bureau
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US)
6. ☒ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☐ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). **UNEXECUTED**
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Item 11. to 16. below concern other document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A FIRST preliminary amendment.
☐ A SECOND or SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information:
Form PCT/IB/308
Copy of International Preliminary Examination Report
Computer Readable Form of Sequence Listing
Statement under 37 CFR 1.821(f)
Executed Small Entity Status Declaration


U.S. APPLICATION NO (if known, see 37 CFR 1.5) 09/554414		INTERNATIONAL APPLICATION NO PCT/CA98/01059		ATTORNEY'S DOCKET NUMBER 2055MC/48896	
17. [X] The following fees are submitted: Basic National Fee (37 CFR 1.492(a)(1)-(5)):				CALCULATIONS PTO USE ONLY	
Search Report has been prepared by the EPO or JPO \$840.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) ... \$670.00 No international preliminary examination fee paid to USPTO (37 CFR 1.482) but international search fee paid to USPTO (37 CFR 1.445(a)(2)) \$760.00 Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$ 970.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(2)-(4) \$96.00					
ENTER APPROPRIATE BASIC FEE AMOUNT =				\$ 840.00	
Surcharge of \$130.00 for furnishing the oath or declaration later than [] 20 [X] 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$ 130.00	
Claims		Number Filed	Number Extra	Rate	
Total Claims		31 -20=	11	X \$18.00	\$ 198.00
Independent Claims		4 - 3=	1	X \$78.00	\$ 78.00
Multiple dependent claims(s) (if applicable)				+ \$260.00	\$
TOTAL OF ABOVE CALCULATIONS =				\$1,246.00	
Reduction by 1/2 for filing by small entity, if applicable. Verified Small Entity statement must also be filed. (Note 37 CFR 1.9, 1.27, 1.28).				\$	
SUBTOTAL =				\$ 623.00	
Processing fee of \$130.00 for furnishing the English translation later than [] 20 [] 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				+	
TOTAL NATIONAL FEE =				\$ 623.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28,3.31). \$40.00 per property +				\$	
TOTAL FEE ENCLOSED =				\$ 623.00	
				Amount to be: \$	
				refunded	
				charged \$	
a. [X] A check in the amount of \$ <u>623.00</u> for the filing fee is enclosed b. [] Please charge my Deposit Account No. _____ in the amount of \$ _____ to cover the above fees. A duplicate copy of this sheet is enclosed. c. [X] The Commissioner is hereby authorized to charge any additional fees, which may be required, or credit any overpayment to Deposit Account No. <u>05-1323</u> . A duplicate copy of this sheet is enclosed.					
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.					
SEND ALL CORRESPONDENCE TO: Evenson, McKeown, Edwards & Lenahan, P.L.L.C. 1200 G Street, N.W., Suite 700 Washington, D.C. 20005 Tel. No. (202) 628-8800 Fax No. (202) 628-8844					
				SIGNATURE  J. D. Evans	
				NAME 26,269	
				REGISTRATION NUMBER May 12, 2000	

Figure 1 illustrates the steps of the proposed algorithm for finding a minimum spanning tree. The process starts with a graph with 6 nodes and 7 edges. The algorithm proceeds by selecting edges in increasing order of weight, ensuring that no cycle is formed. The selected edges are: (1,2) with weight 1, (2,3) with weight 1, (3,4) with weight 1, (4,5) with weight 1, (5,6) with weight 1, (1,3) with weight 2, (2,4) with weight 2, (3,5) with weight 2, (4,6) with weight 2, and (1,4) with weight 3. The final minimum spanning tree has a total weight of 10.

09/554414
426 Rec'd PCT/PTO 1 2 MAY 2000

Attorney Docket: 2055MC/48896
PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: MOSHE SZYF ET AL.

Serial No.: UNASSIGNED

Group Art Unit: UNASSIGNED

Filed: MAY 12, 2000

Examiner: UNASSIGNED

Corresponding to PCT/CA98/01059

November 12, 1998

Title: DNA DEMETHYLASE, THERAPEUTIC AND
DIAGNOSTIC USES THEREOF

PRELIMINARY AMENDMENT

Box PCT

Commissioner for Patents
Washington, D.C. 20231

Sir:

Please enter the following amendments to the claims, prior
to the examination of the application during the U.S. National
Phase.

IN THE CLAIMS:

Claim 4, line 2, change "claims" to --claim-- and delete
"or 3".

Claim 10, line 2, change "claims" to --claim-- and delete
"or 3".

Claim 11, line 2, change "claims" to --claim-- and delete "or 3".

Claim 15, line 1, delete "or 3".

Claim 16, line 1, delete "or 3".

Claim 19, line 2, delete "or 2".

Claim 23, line 1, change "one of claims" to --claim-- and delete "or 22".

Claim 27, line 1, delete "or its cDNA of";
line 2, delete "claim 2".

Claim 28, line 1, delete "or its cDNA";
line 2, delete "of claim 2".

Please add the following new claims:

-- 30. Use of the cDNA of claim 2, for changing the state of differentiation of a cell to allow gene therapy, stem cell selection or cell cloning.

31. Use of the cDNA of claim 2, for inhibiting methylation in cancel cells using vector mediated gene therapy. --


R E M A R K S

Entry of the amendments to the claims before examination of the application is respectfully requested. These claims have been amended to remove multiple dependencies.

If there are any questions regarding this Preliminary Amendment or this application in general, a telephone call to the undersigned would be appreciated since this should expedite the prosecution of the application for all concerned.

Respectfully submitted,

May 12, 2000


J. D. Evans
Registration No. 26,269

EVENSON, McKEOWN, EDWARDS
& LENAHAN, P.L.L.C.
1200 G Street, N.W., Suite 700
Washington, DC 20005
Telephone No.: (202) 628-8800
Facsimile No.: (202) 628-8844

Applicant or Patentee: Moshe Szyf et al.
Serial or Patent No.: _____ Atty. Dkt. No.: 1770-183"US-FC
Filed or Issued: _____
For: DNA DEMETHYLASE, THERAPEUTIC AND DIAGNOSTIC USES THEREOF

**VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS
[(37 CFR 1.9(f) AND 1.27 (d)) - NONPROFIT ORGANIZATION]**

I hereby declare that I am an official empowered to act on behalf of the nonprofit organization identified below:

NAME OF ORGANIZATION McGILL UNIVERSITY
ADDRESS OF ORGANIZATION 845 Sherbrooke Street West
Montréal, Québec, Canada H3A 2T5

TYPE OF ORGANIZATION

- ☒ UNIVERSITY OR OTHER INSTITUTION OF HIGHER EDUCATION
☐ TAX EXEMPT UNDER INTERNAL REVENUE SERVICE CODE [26 USC 501(a) AND 501(c)(3)]
☐ NONPROFIT SCIENTIFIC OR EDUCATIONAL UNDER STATUTE OF STATE OF THE UNITED STATES OF AMERICA

(NAME OF STATE _____)

(CITATION OF STATUTE _____)

- ☐ WOULD QUALIFY AS TAX EXEMPT UNDER INTERNAL REVENUE SERVICE CODE [26 USC 501(a) AND 501 (c)(3)] IF LOCATED IN THE UNITED STATES OF AMERICA

- ☐ WOULD QUALIFY AS NONPROFIT SCIENTIFIC OR EDUCATIONAL UNDER STATUTE OF STATE OF THE UNITED STATES OF AMERICA IF LOCATED IN THE UNITED STATES OF AMERICA

(NAME OF STATE _____)

(CITATION OF STATUTE _____)

I hereby declare that the nonprofit organization identified above qualifies as a nonprofit organization as defined in 37 CFR 1.9 (e) for purposes of paying reduced fees under section 41(a) or (b) of Title 35, United States Code with regard to the invention DNA DEMETHYLASE, THERAPEUTIC AND DIAGNOSTIC USES THEREOF by inventor(s) Moshe Szyf; Sanjoy Bhattacharya; and Shyam Ramchandani described in

- ☐ the specification filed herewith
☐ application serial no. _____, filed _____
☐ patent no. _____, issued _____

I hereby declare that rights under contract or law have been conveyed to and remain with the nonprofit organization with regard to the above-identified invention.

If the rights held by the nonprofit organization are not exclusive, each individual, concern or organization having rights to the invention is listed below* and no rights to the invention are held by any person, other than the inventor who could not qualify as an independent inventor under 37 CFR 1.9(c) or by any concern which would not qualify as a small business concern under 37 CFR 1.9(d) or a nonprofit organization under 37 CFR 1.9(e).

*NOTE: Separate verified statements are required from each named person, concern or organization having rights to the invention averring to their status as small entities. [37 CFR 1.27]

NAME _____
ADDRESS _____

- ☐ INDIVIDUAL ☐ SMALL BUSINESS CONCERN ☐ NONPROFIT ORGANIZATION


NAME _____
ADDRESS _____

- ☐ INDIVIDUAL ☐ SMALL BUSINESS CONCERN ☐ NONPROFIT ORGANIZATION

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. [37 CFR 1.28(b)]

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true, and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

NAME OF PERSON SIGNING Alex Navarre
TITLE IN ORGANIZATION Director - Office of Technology Transfer
ADDRESS OF PERSON SIGNING 845 Sherbrooke Street West, Québec, Canada H3A 2T5

SIGNATURE  DATE 09/05/2000

09/554414

526 Rec'd PCT/PTO 12 MAY 2000

Attorney Docket: 2055MC/48896
PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: MOSHE SZYF ET AL.

Serial No.: UNASSIGNED

Group Art Unit: UNASSIGNED

Filed: MAY 12, 2000

Examiner: UNASSIGNED

Corresponding to PCT/CA98/01059

November 12, 1998

Title: DNA DEMETHYLASE, THERAPEUTIC AND
DIAGNOSTIC USES THEREOF

STATEMENT UNDER 37 C.F.R. 1.821(f)


Assistant Commissioner for Patents
Washington, DC 20231

Sir:

The undersigned hereby states that the contents of the paper copy of the sequence listing in the accompanying U.S. National Stage patent application and the accompanying computer readable copy of the sequence listing are the same.

Respectfully submitted,

May 12, 2000



J. D. Evans
Registration No. 26,269

EVENSON, McKEOWN, EDWARDS
& LENAHA
1200 G Street, N.W., Suite 700
Washington, DC 20005
Telephone No.: (202) 628-8800
Facsimile No.: (202) 628-8844
JDE:dcb

50/PRT

DNA DEMETHYLASE, THERAPEUTIC AND
DIAGNOSTIC USES THEREOF

BACKGROUND OF THE INVENTION

5 (a) Field of the Invention

The invention relates to a novel enzyme, DNA demethylase, therapeutic and diagnostic uses thereof.

(b) Description of Prior Art

Many lines of evidence have established that
10 modification of cytosine moieties residing in the dinucleotide sequence CpG in vertebrate genomes is involved in regulating a number of genome functions such as parental imprinting, X-inactivation, suppression of methylation of ectopic genes and differential gene
15 expression (Szyf, M. (1996) Pharmacol. Ther. 70, 1-37). DNA methylation performs its function of differentially marking genes because the distribution of methylated CpGs is tissue- and site- specific forming a pattern of methylation (Szyf, M. (1996) Pharmacol. Ther. 70, 1-
20 37). It is clear that the pattern of methylation is fashioned by a sequence of methylation and demethylation events (Brandeis, M. et al. (1993) Bioassays 15, 709-713) during development and is maintained in the fully differentiated cell (Razin, A. et al. (1980) Science 210, 604-610). While it was originally suggested
25 that DNA demethylation is accomplished by a passive loss of methyl groups during replication (Razin, A. et al. (1980) Science 210, 604-610), it is now clear that an active process of demethylation occurs in embryonal
30 cells (Frank, D. et al. (1991) Nature 351, 239-241), in differentiating cell lines (Razin, A. et al. (1986) Proc. Natl. Acad. Sci. USA 83, 2827-2831; Szyf, M. et al. (1985) Proc. Natl. Acad. Sci. USA 82, 8090-8094) and in response to estrogen treatment (Saluz, H.P. et al. (1986) Proc. Natl. Acad. Sci. USA 83, 7167-7171).
35 Two modes of demethylation have been documented: site

specific demethylation that coincides in many instances with onset of gene expression of specific genes and a general genome wide demethylation that occurs during early development in vivo during cellular differentiation and in cancer cells (Feinberg, A.P. et al. (1983) Nature 301, 89-92; Razin, A. et al. (1986) Proc. Natl. Acad. Sci. USA 83, 2827-2831). The global demethylation is consistent with the hypothesis that a general demethylase activity which is activated at specific points in development or oncogenesis exists. It has been hypothesized that one mechanism regulating the pattern of methylation is the control of expression of methyltransferase (Szyf, M. (1991) Biochem. Cell Biol. 69, 764-767) and demethylase activities (Szyf, M. (1994) Trends Pharmacol. Sci. 7, 233-238). Although extensive information has been obtained on the enzymatic activity responsible for methylation and the regulation of its expression in the last two decades (Szyf, M. (1996) Pharmacol. Ther. 70, 1-37), the identity of the demethylase has remained a mystery. It is clear however that to fully understand how patterns of methylation are formed and maintained and to determine their role in development, physiology and oncogenesis, one has to identify the demethylase enzyme(s). Two main difficulties have inhibited the identification of this enzyme. First, it is believed that demethylation of a methylated cytosine is chemically highly unlikely since it involves breaking a very stable C-C bond. Second, demethylation occurs at very defined stages in development (Brandeis, M. et al. (1993) Bioassays 15, 709-713) and identifying an adequate tissue source for this enzyme is critical.

Whereas no bona fide demethylase has been identified to date, alternative biochemical mechanisms involving exchange of methylated cytosines with non-

methyated cytosines have been described. One previously proposed mechanism is removal of the methylated base by a glycosylase and its replacement with a non-methylated nucleotide utilizing an "excision-repair" mechanism (Razin, A. et al. (1986) Proc. Natl. Acad. Sci. USA 83, 2827-2831). Glycosylase activities that can remove methylated cytosines from DNA have been demonstrated by Vairapandi and Duker (Vairapandi, M. et al. (1993) Nucl. Acids Res. 21, 5323-5327) and more recently by Jost (Jost, J. P. et al. (1995) J. Biol. Chem. 270, 9734-9739). However it is not clear whether this activity is responsible for the general demethylation observed in cellular differentiation. The fact that the activity identified by Jost acts specifically on hemimethylated sequences (which is not the natural substrate in most cases) and can remove thymidines as well as 5-methylcytosines, supports a repair function for this glycosylase-demethylase (Jost, J. P. et al. (1995) J. Biol. Chem. 270, 9734-9739). An alternative mechanism involving a RNA dependent activity has been recently described by Weiss et al. (Weiss et al., 1996). This proteinase-insensitive RNA dependent activity has been shown to catalyze the excision and replacement of a methylated CpG dinucleotide with a nonmethylated CpG dinucleotide that is contained in a DNA-RNA hybrid molecule (Weiss, A. et al. (1996) Cell 87, 709-718). This activity which was identified in differentiating cells in culture was proposed to be involved in demethylation during development. These previous findings demonstrate that the common accepted model in the filed has been that a *bona fide* demethylase does not exist.

It has been previously proposed that the extensive hypomethylation observed in cancer cells might be a consequence of activation of demethylase activity by

oncogenic pathways (Szyf, M. (1994) Trends Pharmacol. Sci. 7, 233-238; Szyf, M. et al. (1995) J. Biol. Chem. 270, 12690-12696). In accordance with this hypothesis we have shown that ectopic expression of v-Ha-ras had induced demethylation activity in the cells (Szyf, M. et al. (1995) J. Biol. Chem. 270, 12690-12696). Using an assay that directly measures the conversion of 3',³²P labeled methyl dCMP (mdCMP) into dCMP, we have shown that nuclear extracts prepared from P19-Ras transfectants bear high levels of demethylase activity (Szyf, M. et al. (1995) J. Biol. Chem. 270, 12690-12696). Building on this observation, we hypothesized that cancer cell lines were a good source for demethylase. However, it is not evident that Ras expression in p19 cells does reflect the situation in cancer cells. P19 is an embryonic cell and expression of Ras might be differentiating them.

It would be highly desirable to be provided with a *bona fide* DNA demethylase (DNA dMTase) to alter developmental programs for therapeutic and biological use.

SUMMARY OF THE INVENTION

In accordance with the present invention, we demonstrate the purification of a *bona fide* DNA demethylase (DNA dMTase) from a human lung cancer cell line A549, determine its kinetic parameters and substrate specificity. The DNA dMTase activity identified in this study converts methyl-dCMP (mdCMP) residing in the dinucleotide sequence mdCpG into dCMP whereas the methyl group is released as a volatile residue which was identified to be methanol. The activity is purified away from any trace amounts of dCTP, is insensitive to the DNA polymerase inhibitor ddCTP, is not affected by the presence of methyl dCTP (mdCTP) in the

reaction and does not exhibit exonuclease or glycosylase activities. The identification of this new enzyme points out to new directions in our understanding of how DNA methylation patterns are formed and altered.

One aim of the present invention is to provide a *bona fide* DNA demethylase (DNA dMTase).

In accordance with the present invention there is provided a DNA demethylase enzyme having about 40 KDa, and wherein the DNA demethylase enzyme is over-expressed in cancer cells and not in normal cells.

In accordance with the present invention there is provided a cDNA encoding human demethylase which comprises a sequence set forth in SEQ ID NO:1.

In accordance with the present invention there is provided two mouse cDNAs homologous to the human cDNA, wherein the cDNA encoding mouse demethylase having a sequence set forth in SEQ ID NOS:5-7.

In accordance with the present invention there is provided a different human cDNA which encodes a protein homologous to the human demethylase having a sequence set forth in SEQ ID NO:3.

In accordance with the present invention there is provided the use of the expression of demethylase cDNAs to alter DNA methylation patterns of DNA *in vitro* in cells or *in vivo* in humans, animals and in plants.

The demethylase cDNAs expression may be under the direction of mammalian promoters, such as CMV.

The demethylase cDNAs expression may be under plant specific promoters to alter methylation in plants and to allow for altering states of development of plants and expression of foreign genes in plants.

The demethylase cDNAs expression may be in the antisense orientation to inhibit demethylase in cancer cells for therapeutic processes.

5 In accordance with the present invention there is provided the use of the expression of demethylase cDNAs in bacterial or insect cells for production of large amounts of demethylase.

In accordance with the present invention there
15 is provided the use of the sequence of demethylase
cDNAs as a template to design antisense oligonucleo-
tides and ribozymes.

In accordance with the present invention there is provided the use of expression of cDNAs in two hybrid systems in yeast to identify proteins interact-
25 ing with demethylase for diagnostic and therapeutic purposes.

In accordance with the present invention there is provided a volatile assay for high throughput

screening of demethylase inhibitors as therapeutics and anticancer agents which comprises the steps of:

- a) using transcribed and translated demethylase cDNAs in vitro to convert methyl-cytosine present in methylated DNA samples to cytosine present in DNA and volatilize methyl group;
- b) determining the absence or minute amount of volatilize methyl group as an indication of an active demethylase inhibitor.

In accordance with the present invention there is provided a volatile assay for the diagnostics of cancer in a patient sample which comprises the steps of:

- a) determining demethylase activity in patient samples by assaying conversion of methyl-cytosine present in methylated DNA to cytosine present in DNA and its volatilization as methyl groups released as methanol;
- b) determining the presence or minute amount of volatilized methyl released as methanol groups as an indication of cancer in the patient sample.

In accordance with the present invention there is provided the use of an antagonist or inhibitor of DNA demethylase for the manufacture of a medicament for cancer treatment, for restoring an aberrant methylation pattern in a patient DNA, or for changing a methylation pattern in a patient DNA.

Such an antagonist is a double stranded oligonucleotide that inhibits demethylase at a K_i of 50nM, such as $[C^mGC^mGC^mGC^mG] \cdot [G^mCG^mCG^mCG^mC]_n$.

The inhibitors include, without limitation an anti-DNA demethylase antibody, an antisense of DNA demethylase or a small molecule such as any derivative of imidazole.

The change of the methylation pattern may activate a silent gene. Such an activation of a silent gene permits the correction of genetic defect such as found for β -thalassemia or sickle cell anemia.

- 5 The DNA demethylase of the present invention may be used to remove methyl groups on DNA *in vitro* such as needed for cloning DNA.

- 10 The DNA demethylase of the present invention or its cDNAs may be used, for changing the state of differentiation of a cell to allow gene therapy, stem cell selection or cell cloning.

- 15 The DNA demethylase of the present invention or its cDNAs may be used, for inhibiting methylation in cancer cells using vector mediated gene therapy.

- 20 In accordance with the present invention there is provided an assay for the diagnostic of cancer in a patient, which comprises determining the level of expression of DNA demethylase by either RT-PCT, ELISA or volatilization assay of the present invention in a sample from the patient, wherein overexpression of the DNA demethylase is indicative of cancer cells.

BRIEF DESCRIPTION OF THE DRAWINGS

- 25 Figs. 1A to 1B illustrate the purification of demethylase (DNA dMTase) from human A549 cells;

Figs. 2A and 2C illustrate that DNA dMTase is a protein inhibited by RNA and not by ddCTP, mdCTP;

Figs. 2B and 2D illustrate the kinetics of DNA dMTase activity;

- 30 Figs. 3A to 3C illustrate the product of DNA dMTase activity is cytosine and it exhibits no exonuclease or glycosylase activity;

Figs. 4A-4C illustrate the demethylation reaction releases methanol as a volatile residue;

Fig. 4D illustrates the transfer of a proton from water to regenerate cytosine;

Figs. 4E-4F illustrate that the volatile product is methanol;

5 Fig. 5 illustrates the suggested demethylation reaction;

Figs. 6A-6D illustrate the substrate Specificity of DNA dMTase;

10 Figs. 7A-7D illustrate chromatographic isolation of dMTase from human A549 cells;

Figs. 8A-8B illustrate the alignment between the MDB domain of MeCP2 and demethylase and the predicted amino acid sequence of human demethylase;

15 Fig. 8C illustrates the mRNA encoded by demethylase;

Figs. 9A-9F illustrate the cDNA and their predicted amino acid of demethylases and homologues of the present invention (SEQ ID NOS:1-8);

20 Figs. 10A-B illustrate a mammalian expression vector of dMTase and *in vitro* translated dMTase polypeptide;

Fig. 10C illustrates that *in vitro* translated DNA dMTase releases volatile methyl residues from methylated DNA;

25 Fig. 10D illustrates that *in vitro* translated DNA dMTase transform methylated cytosines to cytosines;

Fig. 11A illustrates that transiently transfected demethylase releases volatile residues from methylated DNA;

30 Fig. 11B illustrates the polypeptide expressed from transiently transfected demethylase;

Figs. 11C-11E illustrate that transiently transfected demethylase transforms methylated cytosines to cytosines in a protein dependent manner;

Fig. 11F illustrates that the transformation of methylated cytosine to cytosine by transiently transfected demethylase depends on the concentration of substrate;

5 Fig. 12A illustrates that transiently transfected demethylase catalyzes the transfer of a proton from tritiated water to regenerate cytosine;

Fig. 12B illustrates that the cloned demethylase releases methanol from methylated DNA;

10 Figs. 13A-13C illustrate that the cancer cells express demethylase activity whereas normal cells do not;

Fig. 13D illustrates that demethylase mRNA is highly express in cancer cells;

15 Fig. 14A illustrates demethylase bacterial retroviral and mammalian expression vector;

Fig. 14B illustrates inhibition of demethylase activity by a specific inhibitor;

20 Fig. 14C illustrates inhibition of tumorigenesis *in vitro* by an inhibition of demethylase;

Fig. 15 illustrates inhibition of tumorigenesis in cell culture by induced expression of demethylase antisense vector;

25 Fig. 16 illustrates the inhibition of demethylase by a small molecule inhibitor imidazole; and

Fig. 17 illustrates a model for the inhibition of cancer growth by an inhibition of demethylase.

DETAILED DESCRIPTION OF THE INVENTION

30 The pattern of methylation is fashioned during development by a sequence of methylation and demethylation events. The identity of the demethylase has remained a mystery and alternative biochemical activities have been shown to demethylate DNA but no activity
35 that can truly remove methyl groups from DNA has been


shown to date. Utilizing human lung carcinoma cells as a source for demethylase activity we demonstrate that mammalian cells bear a *bona fide* DNA demethylase (DNA dMTase) activity. DNA dMTase transforms methyl-C to C by catalyzing replacement of the methyl group on the 5 position of C with a hydrogen derived from water. DNA dMTase demethylates both fully methylated and hemimethylated DNA, shows dinucleotide specificity and can demethylate mdCpdG sites in different sequence contexts. This enzyme is different from previously described demethylation activities: it is proteinase sensitive, activated by RNase and releases different products.

DNA dMTase is a novel enzyme showing a new and unexpected activity that has not been previously described in any organism. The finding of a *bona fide* demethylase, points out new directions in our understanding of the biological role of DNA methylation.

In spite of the fact that it was previously shown that Ras expression in p19 cells can induce demethylation activity. It was not clear whether this demethylation activity is indeed a *bona fide* demethylase. One would predict that demethylase is present in embryonal cells. It was surprising to see that demethylation activity is present in cancer cells. The finding of high levels of demethylase in A549 cells is indeed an unexpected discovery.

In accordance with the present invention, it is shown and demonstrated that demethylation occurs by removal of a methyl group from methylated cytosine in DNA, that a hydrogen from water replaces the methyl group at the 5' position, that the resulting methyl group reacts with the remaining hydroxyl from water to generate methanol which volatilizes (Fig. 4E-F). Thus,

bona fide demethylation of DNA involves the following reaction:



- 5 The cDNA cloned in accordance with the present invention is the demethylase since it can convert methyl-cytosines in DNA to cytosines and volatilize the methyl groups on DNA when transcribed and translated in vitro which are released as methanol. This is a novel
- 10 cDNA encoding a biochemical activity that has been not described before.

In accordance with the present invention, there is shown a model for the inhibition of cancer growth by an inhibition of demethylase (Fig. 17).

15

EXPERIMENTAL PROCEDURES

Cell Culture

- A549 Lung Carcinoma cells (ATCC: CCL 185) were grown in Dulbecco's modified Eagle's medium (with low glucose) supplemented with 10% fetal calf serum, 2 mM glutamine, 10 U/ml ciprofloxacin. Human Skin Fibroblasts #72-213A MRHF were obtained from BioWhittaker, Bethesda and were grown in Dulbecco's modified Eagle's medium supplement with 2% fetal calf serum, 2 mM glutamine. H446 Lung carcinoma cells (ATCC: HTB 171) was
- 20 grown in RPMI 1640 medium with 5% fetal calf serum.
- 25

Preparation of nuclear extract

- Nuclear extracts were prepared from A549 cultures at near confluence as previously described (Szyf et al., 1991; Szyf et al., 1995). The cells were trypsinized, collected and washed with phosphate-buffered saline and suspended in buffer A (10 mM Tris, pH 8.0, 1.5 mM MgCl₂, 5mM KCl, 0.5% NP-40) at the concentration of 10⁸ cells per ml for 10 min. at 4°C. Nuclei were
- 30
- 35 collected by centrifugation of the suspension at 1000 g

for 10 minutes. The nuclear pellet was resuspended in buffer A (400 μ l) and collected as described in the experimental procedures. A nuclear extract was prepared from the pelleted nuclei by suspending them in
5 buffer B (20 mM Tris, pH 8.0, 25% glycerol, 0.2 mM EDTA and 0.4 mM NaCl) at the concentration of 3.3×10^8 nuclei per ml and incubating the suspension for 15 min. at 4°C. The nuclear extract was separated from the nuclear pellet by centrifugation at 10,000g for 30 min-
10 utes. Nuclear extract were stored in -80°C for at least two months without loss of activity.

Chromatography on DEAE-Sephadex

A freshly prepared nuclear extract (1 ml , 1.1 mg) was passed through a Microcon™ 100 spin column, the
15 retainant was diluted to a conductivity equivalent to 0.2 M NaCl in buffer L and applied onto a DEAE-Sephadex column (Pharmacia) (1.0 x 5 cm) that was preequilibrated with buffer L (10 mM Tris-HCl, pH 7.5, 10 mM
20 MgCl₂) containing 0.2 M NaCl at a flow rate of 1 ml/min. The column was then washed with 15 ml of the starting buffer (buffer L + 0.2 M NaCl) and proteins were eluted with 5 ml of a linear gradient of NaCl (0.2-5.0 M). 0.8 ml fractions were collected and
25 assayed for demethylase activity after desalting through a Microcon™ 10 spin column (Amicon) and resuspension of the retainant in 0.8 ml buffer L. DNA demethylase eluted between 2-5.0 M NaCl.

Chromatography on S-Sepharose

Active DEAE-Sepharose column fractions were
30 pooled, adjusted to 0.1 M NaCl by dilution and loaded onto an S-sepharose column (Pharmacia) (1.0 x5 cm) which had been preequilibrated with buffer L containing 0.2 M NaCl at a flow rate of 1 ml/min. Following washing of the column as described in experimental proce-
35 dures, the proteins were eluted with 5 ml of a linear

NaCl gradient (0.2-5.0M). 0.5 ml fractions were collected and assayed for DNA demethylase activity after desalting and concentrating to 0.2 ml using a Microcon™ 10 spin column. DNA demethylase activity eluted around 5.0 M NaCl.

Chromatography on Q-Sepharose

Active fractions from S-sepharose column were pooled, adjusted to 0.2 M NaCl by dilution and applied onto a Q-sepharose (Pharmacia) column (1.0 x5 cm) which had been equilibrated as described in the experimental procedures at a flow rate of 1 ml/min. The column was washed and the proteins were eluted with a linear NaCl gradient (0.2- 5.0 M). Fractions (0.5 ml) were collected, assayed for demethylase activity after desalting and concentrating to a final volume of 0.2 ml as described in the experimental procedures. The demethylase activity eluted around 4.8-5.0 M NaCl.

Gel-Exclusion Chromatography on DEAE-Sephacel

The pooled fractions of Q-sepharose column were adjusted to 0.2 M NaCl, loaded onto a 2.0 x 2.0 cm DEAE-Sephacel column (Pharmacia) and eluted with 10 ml of buffer L containing 0.2 M NaCl. The fractions (0.8 ml) were collected and assayed after concentration to about 180 µl with a Microcon™ 10 spin column for DNA demethylase activity. The activity was detected at fraction 4, which is very near the void volume (~200kDa).

Assay of DNA demethylase activity

To directly assay DNA demethylase activity in vitro two independent methods were applied.

(A) To assay the conversion of methyl-dCMP (mdCMP) to dCMP we used a previously described method (Szyf et al., 1995). Briefly, $\alpha^{32}\text{P}$ labeled, fully methylated poly[mdc³²PdG]n substrate was prepared as follows. One hundred ng of a double-stranded fully methylated

(mdCpdG) oligomer (Pharmacia) were denatured by boiling, which was followed by partial annealing at room temperature. The complementary strand was extended with Klenow fragment (Boehringer Mannheim) using methyl-5-dCTP (mdCTP, 0.1 mM) (Boehringer Mannheim) and [α-³²P] GTP (100 μCi, 3000 Ci/mmol), and the unincorporated nucleotides were removed by chromatography through a NAP-5 column (Pharmacia). The NAP-5 chromatography was repeated to exclude minor contamination with unincorporated nucleotides. As a control a non-methylated poly[dC³²pdG]n substrate was similarly prepared except that a nonmethylated dCpdG oligomer served as a template and dCTP was used in the extension reaction. The column fractions (30 μl), described in the experimental procedures were incubated with 1 ng of poly[mdC³²pdG]n substrate for 1 hour at 37°C in a buffer L containing 25% glycerol (v/v) and 5 mM EDTA. The reacted DNA as well as a nonmethylated poly[dC³²pdG]n and methylated [mdC³²pdG]n nonreacted controls were purified by phenol/chloroform extraction and subjected to micrococcal nuclease digestion (100 μg at 10 μl) and calf spleen phosphodiesterase (2μg) (Boehringer) (Pharmacia) to 3' mononucleotides for 15 hours at 37°C. The digestion products were loaded onto a thin layer chromatography plate (TLC) (Kodak, 13255 Cellulose), separated in a medium containing, 132 ml Isobutyric acid:40 ml water: 4 ml ammonia solution, autoradiographed and the intensity of the different spots was determined using a phosphorimager (Fuji, BAS 2000). ³²P labeled substrates and tritium labeled substrates were phosphoimaged using BAS 2000 plate and BAS-TR2040 phosphorimager plate respectively.

(B) The second method determined removal of methylated residues from methylated DNA by measuring disappearance of ³H-CH₃ or ¹⁴C-CH₃ from the reaction mixture. 100 ng

of poly [dCdG]n double stranded DNA was methylated using SssI methylase (New England Biolabs) and an excess of [³H-methyl AdoMet (80 Ci/mmol; New England Nuclear)]. The tritiated methyl group containing DNA was purified from labeled AdoMet using NAP-5 column chromatography. All column purified fractions of DNA demethylase were assayed using the tritiated substrate. In a typical assay, 1 ng of DNA was incubated (at a specific activity of 4 x10⁶dpm/mg) with 30 µl of column fraction for one hour at 37°C in buffer L. To determine the number of methyl groups remaining in the DNA following incubation with the different fractions, 250 µl of water were added and the mixture was incubated at 65°C for 5 minutes. One hundred µl of the reaction mixture were withdrawn for liquid scintillation counting. Controls received similar treatment except that in place of a column fraction, an equal volume of buffer L was added. The number of methyl groups that were removed from the DNA by the different fractions was determined by subtracting the remaining counts in each of the fractions from the counts remaining in the control. All tests were carried out in triplicates. The results are presented as picomole methyl group removed. One unit of DNA dMTase activity is defined as: amount of enzyme that releases one picomole of methyl group from methylated dCp dG substrate in one hour at 37°C.

Methyl removal assay using double-labeled substrates

To determine whether the methyl group leaves the DNA and not any non-specific removal of tritium, we prepared SK plasmid DNA containing a tritiated hydrogen at the 6' position of cytosine and thymidine by growing the plasmid harboring bacteria in the presence of deoxy [6-³H] Uridine (22 Ci/mmol; Amersham) (10µCi/ml). The [6-³H]-cytosine containing pBluescript SK(+) was puri-

fied according to standard protocols and was methylated using an excess of [^{14}C -methyl] AdoMet (59 mCi/mmol; Amersham) (10 μCi per 100 μl reaction) and SssI methylase. The double labeled DNA substrate was purified twice on a NAP-5 column. 15 μl of DNA dMTase were incubated with 1 ng of double labeled DNA (specific activity of 2000 dpm/ng) for 1 hour at 37°C. Following incubation, the remaining ^{14}C versus ^3H counts were determined as described in the experimental procedures by scintillation counting (Wallac). The ^{14}C counts were normalized against ^3H counts. The controls received similar treatment except that instead of DNA dMTase, an equal amount of distilled water was added to them.

To determine the number of $^3\text{H-CH}_3$ in the gaseous phase, 1 ng of $^3\text{H-CH}_3$ poly [dCpdG] DNA were incubated with DNA dMTase overnight in a sealed tube (Pierce, Illinois, USA). 0.8 ml of air were removed from the tube using a gas tight syringe (Hamilton, Reno, Nevada) and injected into a sealed gas tight scintillation vial containing 10 ml OptiPhase scintillation fluid (Wallac, UK) and counted. As a control the DNA was incubated with an equal volume of buffer L and treated similarly.

Synthesis of other methylated dC dinucleotides

Poly [mdC 32 pdA] and [mdC 32 pdT] substrates were prepared as follows. About 0.5 μg of 20 mer oligonucleotides 5'(GG)103', 5'(GT)103' and 5'(GA)103' were boiled and annealed at room temperature with oligonucleotide 5'CCCCC3', 5'CACACA3' and 5'CTCTCT3' respectively. The complementary strand was extended with Klenow fragment using m5dCTP (Boehringer Mannheim) and either [$\alpha^{32}\text{P}$] dATP (100 μCi , 3000 Ci/mmol) or [$\alpha^{32}\text{P}$] dTTP (100 μCi , 3000 Ci/mmol) respectively. The unincorporated nucleotides were removed by chromatography through a NAP-5 column. Hemimethylated mdCpG substrate was prepared in a similar manner except that a nonmeth-

ylated poly dCpdG substrate (Boehringer) was used as template and m5dCTP and [$\alpha^{32}\text{P}$]dGTP were used for extension as described in the experimental procedures.

Assay for nuclease and glycosylase activity

5 [$^{32}\text{pmdCpdG}$]n substrate which included a labeled ^{32}P 5' to mdC was prepared as follows. About 100 ng of poly dCpdG DNA were boiled and partially annealed at room temperature. [$\alpha^{32}\text{P}$]dCTP and cold dGTP were used for complementary strand extension as described in the
10 experimental procedures. The free nucleotides were separated using NAP-5 column chromatography. The purified [$^{32}\text{pmdCpdG}$]n DNA was subjected to methylation by SssI methylase using 320 μM AdoMet. The DNA was repurified twice using a NAP-5 column. The methylated DNA (1
15 ng) was incubated with either 30 μl DNA dMTase, nuclear extract or buffer L. To determine whether $\alpha^{32}\text{P}$ labeled residue is excised from the DNA it was directly applied (3 μl) onto a TLC plate. To determine whether the DNA was demethylated it was subjected to digestion with
20 snake venom phosphodiesterase (0.2 mg in a 10 μl reaction volume) (Boehringer Mannheim) which attacks the 3'-OH group releasing 5'-mononucleotides. The resulting mononucleotides were separated on TLC plates and autoradiographed.

25 To test whether dCTP copurifies with DNA dMTase, which may be involved in activities other than *bona fide* demethylation, 20 μM of dCTP with 1 μl of $\alpha^{32}\text{P}$ labeled dCTP (3000 Ci/mmol) was loaded onto the column with nuclear extract. The ^{32}P counts were measured in
30 the flow through, washes and in the different fractions. About 1.1 million counts were loaded onto the DEAE-Sepharose column and were all recovered up to fraction 8.

35 To determine whether DNA dMTase contains a DNA polymerase activity, DNA demethylase reactions were

performed in presence of 500 μ M of ddCTP (Pharmacia) or 500 μ M of m5dCTP (Boehringer Mannheim) at initial rate conditions.

To determine whether DNA dMTase is sensitive to RNase or Proteinase K treatment, DNA dMTase was pre-treated for 1 h at 56°C with 200 μ g/ml proteinase K (Sigma). A demethylation reaction was carried out with this pretreated fraction in the usual manner using both demethylation assays described in the experimental procedures. To test the effect of RNA digestion on the demethylation reaction, the fractions from different columns were treated with 100 μ g/ml RNase A (Sigma).

Demethylation of pBluescript SK(+) Plasmid

About 4 μ g plasmid pBluescript SK (Stratagene) was subjected to methylation using SssI methylase. The methylated plasmid (4 ng) was incubated with 30 μ l of DNA dMTase Fraction 4 of DEAE-Sephacel column under standard conditions, extracted with phenol: chloroform and precipitated with ethanol. About 1 ng of the plasmid were subjected to digestion with 10 units each of either of the restriction endonucleases EcoRII (GIBCO-BRL), DpnI, HhaI or HpaII (New England Biolabs) before and after methylation as well as after DNA dMTase treatment in a reaction volume of 10 μ l for 2 hour at 37°C. Following restriction digestion the plasmids were extracted with phenol:chloroform, ethanol precipitated and resuspended in 10 μ l. The plasmids were electrophoresed on a 0.8% (w/w) Agarose gel, transferred onto a Hybond Nylon membrane and hybridized with pBluescript SK(+) plasmid which was 32 P labeled by random-priming (Boehringer Mannheim).

Effect of Redox Reagents (NAD, NADH, NADP, NADPH and FeCl₃) on demethylase activity

The reagents were prepared at 100 μ M concentration and added at a final concentration of 10 μ M to a standard methyl removal assay under initial rate condi-

tions as described in the experimental procedures. The methyl removal activity in presence of each of the cofactors was compared to a control DNA dMTase reaction.

5 **Determination of kinetic parameters**

For determination of kinetic parameters, the demethylation reactions were performed using both assays (generation of dCMP and removal of methyl) as described in the experimental procedures except that
10 varying DNA concentrations from 0.1 nM to 2.5 nM were used in a total volume of 50 μ l including 30 μ l of DNA dMTase. Since it has been established by previous experiments that the reaction proceeds for at least 3 hours, the initial velocity of reaction was measured
15 at one hour intervals. The velocity data was collected at each substrate DNA concentration range stated for both assays. The K_m and V_{max} values for DNA demethylase activity were determined from double reciprocal plots of velocity versus substrate concentration.

20 **Measurements of methanol production catalyzed by demethylase by gas chromatography**

Gas chromatography was performed with a VarianTM model 3400 GC equipped with a 30m StabilwaxTM column (0.053 cm i.d.: Restek Corporation). NitrogenTM was
25 used as carrier gas at a flow rate of 32 ml/min, the injector and detector chambers were at 200 and 300°C respectively. The column was maintained at 40°C for 5 minutes after sample injection.

The demethylase reaction was performed in eppendorf tubes kept within sealed scintillation vials with
30 300 μ l of water as aqueous phase (in radioactive trapping experiments this was replaced by 300 μ l of methanol). The demethylase reaction was initiated in buffer L (10 mM $MgCl_2$, 10 mM Tris-HCl pH 8.0) with 500 ng of
35 tritiated SK plasmid (6000 dpm/ μ l) and 100 μ l of demethylase at 37°C. After overnight incubation at 37°C,

the aqueous phase surrounding the eppendorf tube was transferred to a fresh eppendorf tube, 2 μ l of this mixture was injected in the gas chromatography using a gas tight syringe (Hamilton, Reno, Nevada).

5 **Coupled in vitro transcription translation**

The mRNAs encoded by the pcDNA 3.1/His Xpress demethylase constructs described above were transcribed and translated by coupled transcription-translation using PromegaTM TNT reticulocyte lysate kit (according
10 to manufacturer's protocol), 2 μ g of each construct and 40 μ Ci of [³⁵-S]methionine (1,000Ci/mmol, Amersham) in a 50 μ l reaction volume. To purify non labeled in vitro translated demethylase, coupled in vitro transcription and translation was performed as above but in the pres-
15 ence of cold methionine. The translation products were bound to a ProbondTM nickel column (Invitrogen) and demethylase was eluted according to the manufacturer's protocol with increasing concentrations of imidazole. Demethylase is eluted at 350-500mM imidazole. The imi-
20 dazole eluted demethylase was dialyzed and concentrated by lyophilization.

Gas chromatography coupled with Mass spectrometry (GC-MS) Analyses for identification of volatile product of demethylase catalyzed reaction as methanol
25

The demethylation reactions (volume 50 μ l) were run in conical vials having a total internal volume of 350 microlitres. The vials were closed with a teflon-lined screw cap and left at room temperature for 18 h.
30 The vials were cooled in an ice bath, opened and 10 mg of NaCl and 50 microlitres of toluene were added. The vials were frequently shaken over a period of 1 h. The toluene phases were pipetted into clean vials in a manner to rigorously exclude water carry over. Anhydrous
35 sodium sulfate (5 mg) was added to the toluene extracts to remove water, and the toluene phases were pipetted

into autoinjector vials for GC/MS analysis. Aliquots of 3 microlitres were analyzed under the following instrumental conditions: Instrument: Hewlett-Packard 5988A; Column: 30 m x 0.25 mm i.d. fused quartz capillary with 0.25 micron DB-1 liquid phase, programmed after an initial hold for 1 min at 70 deg at 5 deg/min to 80 deg, then ramped ballistically to 280 deg for bake-out for 5 min; Injector and interface temperatures: 250 deg; Helium flow rate 1.5 ml/min; Mass spectrometer: ion source 200 deg, 70 eV electron impact ionization, scanning from m/z 10 to 50 in full scan mode was begun 6 s after injection, and ceased at 1.5 min to avoid acquisition of the intense toluene solvent peak.

Human A549 cells bear a demethylase activity that could be purified away from dCTP and DNA MeTase

The use of an appropriate cellular source and a direct assay for demethylase activity are obviously critical. As we have previously shown that demethylase activity was induced in response to ectopic expression of the Ras oncogene (Szyf et al., 1995) we reasoned that cancer cells might bear high levels of demethylase activity. Based on preliminary studies demonstrating the presence of high levels of demethylase activity in the human lung carcinoma cell line A549, we have chosen this cell line for our further studies and purification steps. Previous studies have used indirect measures such as increased sensitivity to methylation-sensitive restriction enzymes as indicators of demethylase activity (Weiss et al., 1996; Jost et al., 1995). To directly measure the conversion of 5-mdCMP in DNA to dCMP, we have utilized a completely methylated ³²P labeled [mdC³²pdG]_n double stranded oligomer which we had previously described (Szyf et al., 1995). Following incubation with the different fractions, the

DNA is purified and subjected to cleavage with micrococcal nuclease to 3' mononucleotides. The 3' labeled mdCMP and dCMP are separated by thin layer chromatography (TLC) and the conversion of mdCMP to dCMP is directly determined. This assay provides a stringent test for *bona fide* demethylation and discriminates it from previously described 5mCpC replacement activities (Jost et al., 1995; Weiss et al., 1996). The glycosylase-demethylase activity described by Jost et al. (Jost et al., 1995) will require the presence of a ligase activity and an energy source for replacement of mdC with C to be detected by our assay, whereas the demethylase activity described by Weiss et al. will not be detected since it replaces the intact mdC³²pdG dinucleotide with a cold dCpdG without altering its state of methylation (Weiss et al., 1996).

Nuclear extracts were prepared from A549 cells, applied onto a DEAE-Sephadex column, eluted with a linear gradient from 0.2-5.0M NaCl and the fractions were assayed for demethylase (dMTase) activity as described in the experimental procedures. As shown in Fig. 1(A) a clear peak of dMTase activity is eluted at the high salt fraction 10.

Conversion of methylated cytosine to cytosine:
Nuclear extracts prepared from A549 cells (1.1 mg) were passed through an AMICON™ 100 spin column. The retainant (98.56 mg, 0.2 mg/ml) was loaded onto a DEAE-Sephacrose column, the different chromatographic column fractions eluted by a linear NaCl gradient (0.2-5M) were desalted and (30 µl) incubated with 1 ng of [mdC³²pdG]n double stranded oligomer for 1 hour at 37°C, digested to 3' mononucleotides and analyzed on TLC as described in the experimental procedures. Control methylated (ME) and nonmethylated (NM) [dC³²pdG]n substrates were digested to 3' mononucleotides and loaded on the TLC

plate to indicate the expected position of dCMP and mdCMP. The active fraction is indicated by an arrow. This fraction was loaded on S-Sepharose followed by Q-Sepharose and DEAE-Sephacel fractionation.

- 5 The first chromatography step purified the dMTase activity from the bulk of nuclear protein (Fig. 1B) and is a very effective purification step.

DNA dMTase activity as measured by the release of volatile methyl residues. The different column
10 fractions were incubated with lng (4×10^6 dpm/ μ g) of [3 H]-CH₃-[mdCpdG]n oligomer and the release of volatile methyl residues was determined (-) and presented as total dpn). The results are an average of three independent determinations. Protein concentration was
15 determined using the Bio-Rad Bradford kit (-). The elution profile of 20 μ M of [32 P]- α -dCTP incubated with the protein was determined by scintillation counting of the different DEAE fractions (-) and presented as fraction of dCTP loaded on the column.

- 20 To exclude the possibility that the DNA dMTase activity detected in our assay is carried by the DNA MeTase, we assayed the fractions for DNA MeTase activity using a hemimethylated DNA substrate as previously described (Szyf et al., 1991). As observed in Figure
25 1B DNA MeTase activity is detected in the second and third fractions, thus our fractionation separated DNA dMTase away from the DNA MeTase suggesting that they are independent proteins.

There is a remote possibility that the demethylation observed is not a *bona fide* demethylation but
30 a consequence of a glycosylase removal of mC, followed by removal of the remaining deoxyribose-phosphate by AP (apyrimidine) nuclease, repair of the gap catalyzed by DNA polymerase using trace dCTP contained in the frac-
35 tion and ligation of the break with ligase in the pres-

ence of residual ATP. For this hypothesis to be consistent with our data, four independent enzymes and two cofactors have to cofractionate with DNA dMTase. To exclude the possibility that a trace amount of dCTP is bound to DNA dMTase active fraction, we have added 20 μM of ^{32}P labeled dCTP (10×10^6 cpm) to the nuclear extract and determined its elution profile on the DEAE column. Less than background cpm (10 cpm) were detected in the DNA dMTase active fraction suggesting that our first column purifies dCTP away from the DNA dMTase at least 1×10^6 fold (Fig. 1B). If any dCTP is present in the nuclear extract, the remaining concentration after fractionation on DEAE is well below the K_{ms} of the known DNA polymerases. The possibility that dCTP is so tightly bound to the enzyme that it could not be replaced by the exogenous ^{32}P labeled dCTP is very remote since an enzyme using dCTP as substrate must readily exchange dCTP.

The active fraction 10 was further fractionated sequentially on the following columns: S-Sepharose and Q-Sepharose. The DNA dMTase eluted at the high salt fraction from both columns as determined by the [mC 32 pdG]n demethylation assay (Fig. 1A). The ion exchange chromatography was followed by chromatography on DEAE-Sephacel.

The fact that we have maintained our activity even after 4 fractionation steps (Table 1) and that only a single polypeptide is apparent after the last purification step argues strongly against the possibility that the activity detected in our study is a repair or replacement activity. Any replacement mechanism must involve a number of proteins and additional cofactors and substrates. In summary, the chromatography of the demethylase activity in A459 cells provides strong

support to the hypothesis that mammalian cells bear a *bona fide* demethylase activity.

DNA dMTase releases a volatile derivative

A *bona fide* demethylation has to result in
5 release of the methyl group as a volatile derivative
such as CO₂, methanol, methane or formaldehyde. We
have therefore incubated a {[³H] -CH₃-dCpdG}n double
stranded oligonucleotide with the different column
fractions and the rate of release of the tritiated
10 methyl from the aqueous phase was determined by scin-
tillation counting of the remaining radioactivity in
the reaction mix. As demonstrated in Fig. 1b (dia-
mond), the dMTase active fractions release labeled
methyl groups from the methylated substrate.

15 **DNA dMTase is a protein which is inhibited by RNA, does
not involve an exchange activity and does not require
additional cofactors**

DNA dMTase activity measured either as transfor-
20 mation of mdC to C (Fig. 2a) or as release of volatile
methyl residues (Fig. 2c) is abolished after proteinase
K treatment and is not inhibited but rather enhanced
following RNase treatment. 500 μM of ddCTP which
inhibits DNA polymerase does not inhibit demeth-
25 ylation of the [mdC32pdG]n substrate, nor is it inhib-
ited by high concentrations of methyl-dCTP (500 μM)
(Fig. 2a), which is consistent with the hypothesis that
demethylation does not involve an excision and replace-
ment mechanism. If a replacement mechanism is involved
30 in demethylation, the presence of mdCTP should result
in incorporation of methylated cytosines and essential
inhibition of demethylation. Thus, the DNA dMTase
identified here is a protein and not an RNA and is une-
quivocally different from the previously published RNA
35 based or glycosylase based demethylase activities.

The DNA dMTase reaction proceeds without any requirement for additional substrates such as dCTP, redox factors such as NADH and NADPH or energy sources such as ATP (data not shown). As observed in Fig. 2b and 2d, the DNA dMTase reaction maintains its initial velocity up to 90 minutes and continues up to 120 minutes. This time course is inconsistent with dependence on enzyme-bound additional nonreplenishable substrates such as dCTP or ATP or a nonreplenishable redox factor such as NADH or NADPH. Exhausting the nonreplenishable substrate or redox factor would have resulted in rapid deceleration of the initial velocity.

A product of the demethylation reaction is deoxyCytosine in DNA

What is the product of the demethylation reaction? The results presented above (Fig. 1a, 2a and b) based on a one dimension TLC separation show that DNA dMTase generates dC from mdC in DNA. To further substantiate this conclusion, we subjected DNA dMTase treated DNA to remethylation with the CpG MeTase *M.Sss* I which can transfer a methyl group exclusively to dC. The results presented in Fig. 3a show that the demethylated product of DNA dMTase is dC since it is completely remethylated with *M.Sss* I. The identity of the demethylated product as dC was further established by a two-dimension TLC analysis demonstrating that the product of dMTase comigrates with a cold dCMP standard in both dimensions (Fig. 3b).

DNA dMTase does not release a nucleotide, a phosphorylated base or phosphate from methylated DNA when incubated with a [32pmdCpdG]n substrate which included a labeled 32P 5' to mdC or our standard methylated substrate (Fig. 1) where 32P is 3' to the m5dC (Fig. 3c). Nuclear extracts which obviously contain a number of glycosylases and nucleases release phospho-

rylated derivatives in the same assay (Fig. 3c). dMTase transforms the methyl cytosine in the [32pmdCpdG]n substrate to cytosine as demonstrated when the reacted DNA is digested to 5' mononucleotides (Fig. 3c +V PDS) and analyzed by TLC. Since this reaction does not involve release of a 32P derivative (Fig. 3c -V PDS), it demonstrates that dMTase transforms methylated cytosines to cytosines on DNA without disrupting the integrity of the DNA substrate by glycosylase or nuclease activity .

The second product of the dMTase reaction is methanol

What is the identity of the leaving group? The results presented in Fig1b suggest that the labeled methyl leaves the DNA as a volatile compound. The demethylase reaction involves release of the methyl group *per se* whereas the cytosine base ring remains in the aqueous phase. Fig. 4a demonstrates this point by using a methylated plasmid labeled with a ³H-hydrogen at the sixth position of cytosine and [14C]-methyl at the fifth position of cytosine as a substrate.

The three most obvious candidates the methyl group is leaving as are formaldehyde, carbon dioxide, and methanol. Methadone trapping for labeled formaldehyde detection and sodium hydroxide trapping for labeled carbon dioxide detection were both negative in identifying the form in which the methyl group is leaving in the dMTase reaction (data not shown). The other possible chemical form that the methyl group may leave the DNA as, is methanol. Since methanol is a volatile compound, a simple method to measure generation of methanol is a scintillation-volatilization assay (see Fig. 4b for description). Volatilization assays have been previously used to measure release of methanol in demethylation reactions. The demethylation reaction mix containing the labeled {[³H] -CH₃-dCpdG]n substrate

with either dMTase or no enzyme, as a control, is added to an uncapped 0.5 ml tube which is placed in a sealed scintillation vial containing scintillation fluid. Released methanol is volatile, diffuses out of the open
5 reaction tube and is mixed with the excess of the scintillation fluid in the vial registering as counts in the scintillation counter. As a control indicating that methanol is volatilized under the conditions of our assay, we incubated approximately equal counts of
10 radioactively labeled methanol under the same conditions and measured the counts in a scintillation counter at different time points. As observed in Fig. 4c the majority of methanol in the reaction tube volatilizes from the reaction tube into the scintillation
15 fluid following an overnight incubation at 37°C. The experiment shown in Fig. 4b demonstrates that volatilized label is released from methylated DNA only in the presence of dMTase.

The identity of the volatile group has been
20 determined to be methanol by a gas chromatography (GC) analysis. The demethylation and control reactions (indicated in Fig. 4e) were performed in an uncapped tube placed in a sealed scintillation vial containing a larger volume (300 μ l) of water. The volatile residue
25 diffuses into the surrounding water and mixes with it. A 2 μ l sample of the surrounding water was injected into a GC column as described in the methods. As shown in Fig. 4e, the volatile compound released by dMTase in a dose response manner coelutes with methanol. Release of methanol is observed only in the pres-
30 ence of both dMTase and methylated DNA. No methanol is released when dMTase is reacted with nonmethylated DNA, demonstrating that methanol is a product of demethylation of DNA.

The leaving group was also identified as methanol using gas chromatography coupled with Mass spectrometry (GC-MS). As illustrated in Fig. 4f., incubation of methylated DNA with dMTase (dMTase+ME-DNA) results in release of a peak with the retention time and mass spectrum (peaks are identified at 32 and 29 atomic mass which are the atomic masses of methanol and ionized methanol respectively) which is consistent with its identification as methanol. Incubation of dMTase with nonmethylated DNA does not release methanol indicating that methanol is a product of the demethylation reaction. No methanol is released when the samples are incubated with dMTase treated with protease K indicating that the release of methanol from methylated DNA is catalyzed by an enzymatic activity.

Demethylation involves transfer of a hydrogen from water to regenerate cytosine

If demethylation involves removal of the methyl moiety from mdC, a hydrogen has to be transferred to the carbon at the 5' position to regenerate cytosine. Since no redox factors are involved, what is the source of the hydrogen? To test the hypothesis that the source of the hydrogen is water, we incubated either non labeled [mdCpdG]n or [dCpdG]n double stranded DNA with DNA dMTase for different time periods in the presence of tritiated water, following which the DNAs were digested to 3' dNMPs, separated on TLC with non-radioactive standards for each of the 5 possible dNMPs and exposed to a tritium sensitive phosphorimaging plate. As seen in Fig.4d, dMTase catalyzes the transfer of a tritiated hydrogen from water to dCMP in methylated DNA in a time dependent manner only when methylated DNA is used as a substrate. Based on the experiments described in Fig.3 and 4 we propose that dMTase catalyzes the exchange of the methyl group at

the 5' position of cytosine in DNA with hydrogen from water and the methyl group reacts with the remaining hydroxyl group to form methanol (Fig. 5).

5 Substrate and sequence specificity of DNA dMTase

Methylation of CpG dinucleotides is the most characterized modification occurring in genomic DNA^{8,48}. The results presented in Fig.6 demonstrate that DNA dMTase is a general DNA dMTase activity that demethylates fully or hemimethylated dCpdG in DNA flanked by a variety of sequences which are distributed at different frequencies, but does not demethylate methylated adenines or methylated cytosines that do not reside in the dinucleotide CG. First, as shown in Fig.6a, a plasmid DNA methylated *in vitro* at all dCpdG sites with *M.Sss* I and all d*CdCdGdG sites with *M. Msp* I (which methylates the external C in the sequence *CCGG, thus enabling the determination of demethylation at the CC dinucleotide) and *in vivo* with the *E. coli* DCM MeTase at dCmdCdA/dTdGdG sites and with the DAM MeTase at dGmdAdTdC sites (adenine methylated) was treated with dMTase and the state of methylation of the plasmid was determined using the indicated methylation sensitive restriction enzymes. dMTase demethylates C*G methylated sites as indicated by the sensitivity of the dMTase treated plasmid to *Hpa* II and *Hha* I but does not demethylate C*C,C*A or C*T methylated sites as indicated by the resistance to *Msp* I and *Eco* RII restriction enzymes, or adenine methylation as indicated by its sensitivity to *Dpn* I. Second, bisulfite mapping analysis of methylation of 5 methylated C*G sites residing in a *M.Sss* I *in vitro* methylated pMetCAT plasmid following dMTase treatment shows that all C*G sites are demethylated irrespective of their flanking sequences thus excluding the possibility that demethylation is limited to CCGG or CGCG sequences (Fig. 6b).

Third, dMTase does not demethylate two fully methylated cytosine bearing oligomers [dmC32pdA]n, [mdC32pdT]n demonstrating that mdCpdA and mdCpdT are not demethylated by DNA dMTase (Fig. 6d). Fourth, dMTase demethylates a hemimethylated synthetic substrate [dCpdG]n*[mdC32pdG]n (Fig. 6d). Demethylation of SK is complete under these conditions (Fig. 6a) whereas demethylation of a methylated [mdCpdG]n substrate is not complete under the same conditions (Fig. 6d). This can reflect differences in the sequence composition of the substrate and the frequency of methylated cytosines. The [mdCpdG]n contains on average 16 fold more methylated cytosines per molecule than plasmid DNA. Alternatively, these differences might reflect discrepancies in the assays used, restriction enzyme digestion versus a nearest neighbor analysis. To address this discrepancy we have labeled a fully methylated SK plasmid with [α^{32} P]dCTP, 5-methyl-dCTP and the other dNTPs, subjected it to dMTase treatment and digested it to mononucleotides at different time points following the initiation of the reaction and subjected the samples to a TLC analysis. As shown in Fig. 6c, the SK plasmid is fully demethylated at 3 hours which is consistent with the results obtained with methylation sensitive restriction enzymes (Fig. 6a).

The K_m of DNA dMTase for hemimethylated and fully methylated DNA was determined by measuring the initial velocity of the reaction at different concentrations of substrate (Table 2). The calculated K_m for hemimethylated DNA is 6 nM which is two fold higher than the K_m for DNA methylated on both strands, 2.5-3 nM (Table 2). It is unclear yet whether this small difference in affinity to the substrate has any significance in a cellular context. Thus similar to the DNA MeTase DNA dMTase shows dinucleotide sequence

selectivity but in difference from DNA MeTase which shows preference to hemimethylated substrates dMTase prefers fully methylated DNA which is consistent with a role for DNA dMTase in altering established methylation patterns.

Table 1

Purification of DNA dMTase

Purification step	Total protein (µg)	Total dpm	pMole/µg	pMole/µg/h	Fold Purification
Nuclear extract	6000	1107.2	5.5×10^{-5}	1.833×10^{-5}	-
DEAE-Sephadex	3.75	5844	0.4674	0.156	8445.5
SP-Sepharose	0.77	5106	1.989	0.663	35939.84
Q-Sepharose	0.46	5335	3.4	1.13	62860.65
DEAE-Sephacel	0.018	1834	30.57	10.19	552243.2

10

Table 2

Kinetic parameters for DNA dMTase

Method	K_m (DNA)	V_{max} (pMole/h)
Methylated oligo CpG	2.5 nM	340
Hemi-methylated CpG	6.0 nM	402
Methylated SK-DNA	3.3 nM	40.42

Cloning and construction of demethylase expression vectors

15 PCR amplification of the MBD domain of the putative demethylase candidate cDNA

One µg of total RNA prepared from the human small lung carcinoma cell line A549 was reverse transcribed using Superscript reverse transcriptase and random primers (Boehringer) in a 25 µl reaction volume according to conditions recommended by the manufacturer (GIBCO-BRL). Five µl of reverse transcribed cDNA were subjected to an amplification reaction with Taq polymerase (Promega, 1 unit) using the following set of

primers: sense 5'CTGGCAAGAGCGATGTC 3' SEQ ID NO:9,
antisense 5'AGTCTGGTTTACCCTTATTTTG 3' SEQ ID NO:10.

Amplification conditions were: step 1. 95°C 1 min.; step 2: 94°C 0.5 min; step 3: 45°C 0.5 min.; step 4: 72°C 1.5 min; steps 2-4 were repeated 30 times. MgCl₂ was adjusted to 1 mM according to conditions recommended by the manufacturer. The PCR products were cloned in pCR2.1 vector (Invitrogen) and the sequence of the cDNAs was verified by dideoxy-chain termination method using a T7 DNA sequencing kit (Pharmacia). The amplified fragment was excised from the plasmid with EcoRI, labeled with a Boehringer random prime labeling kit according to manufacturer's protocol and alpha ³²P-dCTP. The labeled probe was used to screen a HeLa cell cDNA library in λTriplEx phage (Clontech) according to standard procedures. Positive clones were identified and further purified by serial dilutions for 4 rounds. The insert in the pTriplEx plasmid was excised from the phage according to manufacturer's protocols and the identity of the insert was verified by sequencing. The insert was excised by NotI restriction and subcloned into either the inducible expression vector: Retro tet on (Clontech) in the sense and antisense orientation or the pCDNA3.1/His Xpress vector in all three frames and in the antisense orientation.

Transfection and expression of demethylase in vertebrate cells

Ten μg of either Retro tet on demethylase or pCDNA 3.1/His Xpress demethylase are mixed with 8 μl of transfection lypophilic reagent Pfx-2 (Invitrogen) and placed upon 100,000 mouse (3T3 Balb/c, human (A549) or monkey cells (CV-1) according to manufacturer's protocol in OPTIMEM medium for 4 hours. Cells are harvested after 48 hours and demethylation and demethylase activity is determined by measuring total genomic DNA meth-

ylation using standard techniques or a cotransfected in vitro methylated plasmid using a HpaII /MspI restriction enzyme analysis. Cellular transformation is measured by a soft agar assay.

5

Demethylation of pBluescript SK(+) Plasmid

About 4 μ g plasmid pBluescript SK (Stratagene) was subjected to methylation using SssI methylase. The methylated plasmid (4 ng) was incubated for different time points as indicated with 30 μ l of DNA dMTase Fraction 4 of DEAE-Sephacel™ column under standard conditions, extracted with phenol: chloroform and precipitated with ethanol. About 1 ng of the plasmid were subjected to digestion with 10 units each of either of the restriction endonuclease EcoRII (GIBCO-BRL), DpnI, or HpaII (New England Biolabs) before and after methylation as well as after DNA dMTase treatment in a reaction volume of 10 μ l for 2 hour at 37°C. Following restriction digestion the plasmids were extracted with phenol:chloroform, ethanol precipitated and resuspended in 10 μ l. The plasmids were electrophoresed on a 0.8% (w/w) Agarose gel, transferred onto a Hybond™ Nylon membrane and hybridized with pBluescript SK(+) plasmid which was ³²P labeled by random-priming (Boehringer Mannheim).

dMTase activity coelutes with a ~45 KDa polypeptide when sized under denaturing conditions but migrates as a higher molecular weight complex under non denaturing conditions. dMTase was purified up to 500,000 fold by four chromatographic steps (Table 1). We first determined the identity of the polypeptide associated with dMTase activity by SDS-PAGE analysis of the active fractions. As observed in Fig. 7a, a cluster of 4 polypeptide bands from ~44 KDa to 35 KDa coelute with dMTase activity in the last two chromatographic steps

(the lower fragment might be a degradation product as evidenced by its abundance in the later chromatographic steps). However when the active DEAE-Sephacel fraction is size fractionated on a 4% non denaturing acrylamide column, the dMTase activity elutes at the high molecular weight of ~170 kDa (Fig. 7c, fraction 63). SDS-PAGE analysis of this fraction (63) reveals only two bands (Fig. 7b) observed in the active chromatographic fractions (Fig. 7a). To further determine whether dMTase is found in a multimeric complex, fraction 63 was size fractionated on a glycerol gradient (Fig. 7d) and DNA dMTase activity eluted at the ~170 kDa range. As only two main small polypeptides were identified in fraction 63 (approximately 35-43 kDa), dMTase is probably found in either a homomeric complex if only one of the two peptides is dMTase or a heteromeric complex if both polypeptides are associated with dMTase activity.

a. Identification of a lead DNA dMTase candidate by homology search of dbEST

As the purification of dMTase suggests that the dMTase is of very low abundance, only ~19 ng of dMTase could be isolated from 6 mg of nuclear extract (Table 1), we opted for cloning the dMTase based on its following functional properties. First, since dMTase specifically demethylates methylated CG dinucleotides, we assumed that it should bear the ability to recognize methylated CG dinucleotides. Second, the demethylase transforms methylated cytosine in DNA to cytosine. Third, the demethylase releases the methyl group as a volatile compound.

Previous reports have shown that proteins interacting with methylated DNA share a common domain (MDBD). A TBLASTN search of the dbEST database identified a novel expression tag cDNA (from a T-cell lymphoma Homo sapiens cDNA 5' end) (gb/AA361957/AA361957

- EST71295) and the mouse homologue ((gb/W97165/W97165
mf90g05.r1) from Soares mouse embryo NbME13.5) with
unknown function that bears homology to the MDBD
(Fig. 8a). A search of the GenBank database verified
5 that it is a novel cDNA that has not been included in
GenBank. Alignment of the novel EST and MeCP2 and
MeCP1 associated protein has revealed no homology
beyond the previously characterized MDBD which is con-
sistent with a different function for this methylated
10 DNA binding protein. A 201bp fragment bearing the
sequence identified in the search was reverse tran-
scribed and amplified from human lung cancer cell line
A549 RNA and was used to screen a cDNA library from
Hela cells. The largest insert cloned was of 1.36 kb
15 size and its sequence identity with the EST sequence
was determined. The cDNA is novel and has no homologue
in GenBank and no function has ever been assigned to
it. A virtual translation of the protein identified an
open reading frame (ORF) of 262 amino acids (Fig. 8b).
20 The ORF may extend further 5' as no in frame stop codon
was found upstream of this ATG. However, RACE analy-
ses and further searches of the dbEST have failed to
identify 5' sequences upstream to the one identified in
our screening.
- 25 A BLAST search of the candidate protein using
the Predict protein server against a database of pro-
tein domain families has identified only the MDBD
domain and found no homologue to the sequence in the
data base search. No other functional motifs were
30 identified by the Prosite analysis. This is consistent
with a novel biochemical function for this protein. A
coiled coil prediction of the sequence identified a
coiled coil domain which is known to play a role in
protein protein interactions.

The identified cDNA encodes an mRNA that is widely expressed in human cells as revealed by a Northern blot analysis of human poly A+ mRNA (Fig. 8c) as one major transcript of ~ 1.6 kb which is close to the size of the cloned cDNA, verifying that the cloned cDNA does not represent a highly repetitive RNA but rather a mRNA encoded by a single or low copy number gene.

In vitro translated candidate cDNA bears dMTase activity

A conclusive proof for the existence of a single protein that *bona fide* demethylates DNA is to demonstrate that an *in vitro* translated candidate cDNA can volatilize methyl groups from methylated DNA and transform a methyl cytosine to cytosine in an isolated system. The candidate dMTase cDNA was subcloned it into a pcDNA3.1/His Xpress (INVITROGEN) expression vector in the putative translation frame (pcDNA3.1His A) and in a single base frame shift (pcDNA3.1His B), and was *in vitro* transcribed and translated in the presence of ³⁵S-methionine and the resulting translation products were resolved by SDS-PAGE. Autoradiography revealed a ~40KDa protein (Fig. 10a). The apparent size of the *in vitro* translated protein is shorter by ~3-5 KDa from the apparent size of the purified protein. The cloned cDNA might be missing some upstream amino acids as discussed above or might be differently modified in human cells.

Two tests established whether the *in vitro* translated candidate cDNA is a *bona fide* dMTase. We first tested whether *in vitro* translated protein (purified on a Ni²⁺ charged agarose resin) can volatilize and release methyl residues in [³H]-CH₃-DNA using a radioactive trapping volatilization assay. To verify that the volatilized counts are true ³H counts, a spectrum analysis was performed. As demonstrated in Fig.

10b no volatilization of tritiated methyl residues is observed in the misframe dMTase (misframe) whereas in vitro translated putative dMTase cDNA catalyzes the volatilization of $^3\text{H}-\text{CH}_3$ residues which are trapped in the scintillation cocktail.

Second, *in vitro* translated dMTase cDNA transforms CH_3 -cytosine residing in [^{32}P]- α -dGTP labeled plasmid DNA or in [methyl-dC32pdG] $_n$ double stranded oligomer DNA to cytosine, whereas a frame shift in *in vitro* translated dMTase does not demethylate DNA (Fig. 10d). This demonstrates that the dMTase activity is dependent on the dMTase translation product and not a contaminating activity found in the *in vitro* translation kit that copurifies with the putative dMTase. The reaction carried out by the *in vitro* translated dMTase displays: dependence on the dose of *in vitro* translated product (Fig. 10c), time dependence (Fig. 10d) and dependence on translated protein (Fig. 10b & d misframe, Fig. 10c protease K treatment). Taken together, these results strongly suggest that the cDNA cloned here codes for a *bona fide* enzymatic DNA demethylase activity.

Transiently transfected dMTase cDNA demethylates DNA

dMTase cDNA and the pcDNA3.1HisC vector control were transiently transfected into human embryonal kidney cells to test whether the cDNA can direct expression of dMTase activity in human cells. The His-tagged proteins were bound to Ni^{2+} agarose resin and eluted from the resin with increasing concentrations of imidazole. The expression of the transfected dMTase was verified by a Western blot analysis (Fig. 11b). The imidazole fractions were assayed for their ability to volatilize and release methyl residues in [^3H]- CH_3 -DNA using a radioactive trapping volatilization assay 1. As observed in Fig. 11a, imidazole fractions from

dMTase transfected cells volatilize [^3H]- CH_3 , whereas no tritiated counts are detected in DNA treated with imidazole fractions from cells transfected with a misframe mutation of dMTase or non transfected cells. The transiently expressed dMTase transforms methylated cytosine in DNA to cytosine residing in two different substrates (Figs. 11c & 11d), in a protein dependent manner (Figs. 11c & 11e), and the reaction displays substrate dependence and saturability (Fig. 11f). Transiently expressed dMTase was loaded on a non denaturing glycerol gradient to determine its native MW. Similar to dMTase purified from human cells, cloned and purified dMTase activity fractionated at the 160-190 KDa range (data not shown). This is consistent with self association of cloned dMTase possibly mediated by the coiled-coil domain.

Cloned DNA dMTase catalyzes a hydrolysis of 5-methylcytosine to release methanol

We determined the mechanism by which methyl residues are released by the cloned dMTase (from Fig. 11) and compared it to the purified *bona fide* dMTase activity. Increasing amounts of non labeled [methyl-dCpdG] DNA were incubated with either the *bona fide* dMTase activity purified from A549 cells or the cloned dMTase in the presence of [^3H] water for 3 hours followed by digestion to mononucleotides, a thin layer chromatography and autoradiography. As Fig. 12a shows, both reactions replace the methyl group in 5-methylcytosine with a proton donated from water as indicated by the presence of [^3H] label in cytosine.

The identity of the leaving methyl group in the demethylation reaction catalyzed by the purified *bona fide* dMTase activity was shown to be methanol. In order to identify the form that the methyl residue leaves as in the demethylation reaction catalyzed by

the cloned dMTase an identical gas chromatography/mass spectrometry analysis of the reaction products was performed as in 1. Only the properly translated form of dMTase (both in vitro translated and transiently transfected and purified) is able to produce ions characteristic of methanol in a mass spectrometric analysis (mass of 32 and 29, Fig. 12b). These results suggest that the demethylation reaction catalyzed by the cloned dMTase is hydrolysis of the 5-methyl-cytosine to cytosine and methanol as described for the purified dMTase1.

DNA dMTase activity is undetectable in nontransformed cells

The assays for dMTase activity described here and the cloning of DNA dMTase cDNA enables a study of its expression at different cellular states. Global hypomethylation of DNA is a common observation in cancer cells. This has been a perplexing observation, since DNA MeTase activity is elevated in cancer cells. Hyperactivation of DNA MeTase has been proposed to play a role in cancer development. This paradox raises questions on the proposed role of the elevated levels of DNA MeTase in cancer cells. One simple explanation that has been previously suggested to resolve this paradox is that cancer cells express induced levels of DNA dMTase. We compared the DNA dMTase activity in equal concentrations of DEAE-Sephadex fractionated nuclear extracts (fractions 9-10) prepared from a number of carcinoma cell lines H446, Colo 205, Hela, and A549 with a similar preparation from human skin fibroblast cells at initial rate conditions using [mdC32pdG]n double stranded oligomer as a substrate. As observed in Fig. 13a, whereas DNA dMTase activity is readily observed in all carcinoma cell lines, it is undetectable in nontransformed human cells. The absence of dMTase activity in human primary cells

reflects the situation *in vivo* since dMTase activity is undetectable in preparations from different murine tissues whereas dMTase activity is present in a murine carcinoma cell line P19 that was transfected with the H-Ras protooncogene, or human tumors carried as xenografts in the same strain of mouse (Fig. 1a: COLO 205, A549, Hela). These conclusions were verified using the radioactive-trapping volatilization assay shown in Fig. 13c.

Since dMTase mRNA has been detected using a sensitive poly A+ Northern blot in all normal human tissues, we tested the hypothesis that the absence of detected dMTase activity in normal tissues reflects a quantitative difference in DNA dMTase mRNA between normal tissues and cancer lines. A Northern blot analysis and quantification of dMTase mRNA by a slot blot analysis shown in Fig. 13d using total RNA supports this hypothesis. Whereas minute levels of dMTase mRNA are detected in normal tissues, high levels of dMTase are expressed in a murine carcinoma cell line Y1 that bears a 30 fold amplification of *Ha-ras*.

A second DNA demethylase dMTase2 identified in human and mouse

cDNA sequences, predicted amino acid sequences, and GenBank accession numbers of both dMTase1 and dMTase2 from human and mouse are shown. We claim that the high level of identity of the two proteins (Figs 9c and e) suggests that the two proteins can perform the same function, DNA demethylation. The N-terminals of dMTase1 and dMTase2 contain a Methylated DNA Binding Domain (MBD) and near their C-terminals is a coiled-coil domain, however the middle portions of the protein sequences have no homology to any known structural or catalytic motif. Importantly, their middle regions are still extensively homologous suggesting that the cata-

lytic site of the demethylase activity lies in this area on both proteins.

Induced expression of DNA demethylase in the Antisense orientation inhibits tumorigenesis ex vivo

- 5 To test the hypothesis that inhibition of DNA dMTase can inhibit tumorigenesis tetracycline inducible vectors carrying the human dMTase1 cDNA in either the sense or antisense orientation were constructed and transiently transfected into HEK 293 cells, treated for 10 48 hours either in the presence or absence of doxycycline (a tetracycline analogue), selected for the last 24 hours with puromycin, and then plated on soft agar and allowed to grow for seven days. After seven days colonies were scored and the data presented clearly 15 show that doxycycline induced expression of the dMTase1 cDNA in the antisense orientation reduced colony formation (Fig. 15).

Imidazole is a small molecule inhibitor of DNA demethylase activity

- 20 A template small molecule, imidazole, was tested for the ability to inhibit DNA dMTase activity. In a volatilization of radioactive methyl residues assay, concentrations from 1 μ M to 10mM of imidazole were incubated in a typical volatilization of radioactive methyl 25 residues as described above. The graph clearly demonstrates a dose dependent inhibition of DNA dMTase activity by imidazole, and validates a rationale for testing imidazole based molecules as inhibitors of DNA dMTase activity (Fig. 16).

- 30 **Identification of DNA demethylase cDNAs and protein sequences**

- Fig. 9a** illustrates cDNA sequence of human dMTase1 (SEQ ID NO:1) and its predicted amino acid sequence (SEQ ID NO:2), including its Genbank location. **Fig. 9b** illustrates cDNA sequence of human dMTase2 (SEQ ID NO:3) and its predicted amino acid sequence (SEQ ID NO:4), includ-
- 35

ing its GenBank location. **Fig. 9c** illustrates protein sequence alignment of human dMTase1 and human dMTase2. **Fig. 9d** illustrates cDNA sequence of mouse dMTase1 (SEQ ID NO:5) and its predicted amino acid sequence (SEQ ID NO:6), including its GenBank location. **Fig. 9e** illustrates cDNA sequence of mouse dMTase2 (SEQ ID NO:7) and its predicted amino acid sequence (SEQ ID NO:8), including its GenBank location. **Fig. 9f** illustrates protein sequence alignment of mouse dMTase1 and mouse dMTase2.

While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth, and as follows in the scope of the appended claims.

ART 34 AMDT

WHAT IS CLAIMED IS:

1. A DNA demethylase enzyme and/or homologue thereof, which comprises an amino acid sequence set forth in SEQ ID NOS:2, 4, 6 and 8 and homologue thereof, and wherein said DNA demethylase enzyme has about 40 KDa and is overexpressed in cancer cells.
2. A cDNA encoding a human demethylase of claim 1, which comprises a sequence set forth in SEQ ID NOS:1 and 3.
3. A cDNA homologous to the cDNA of claim 2, wherein said cDNA encoding mouse demethylase set forth in SEQ ID NOS:5 and 7.
4. The use of the expression of demethylase cDNA of claims 2 or 3 to alter DNA methylation patterns of DNA in vitro in cells or in vivo in humans, animals and in plants.
5. The use of claim 4, wherein said demethylase cDNA expression is under the direction of mammalian promoters.
6. The use of claim 5, wherein said promoter is CMV.
7. The use of claim 4, wherein said demethylase cDNA expression is under plant specific promoters to alter methylation in plants and to allow for altering states of development of plants and expression of foreign genes in plants.
8. The use of claim 4, wherein said demethylase cDNA expression is in the antisense orientation to inhibit demethylase in cancer cells for therapeutic processes.

9. The use of claim 9, wherein expression of demethylase cDNA in mammalian cells is to alter their differentiation state and to generate stem cells for therapeutics, cells for animal cloning and to improve expression of foreign genes.

10. The use of the expression of demethylase cDNA of claims 2 or 3 in bacterial or insect cells for production of large amounts of demethylase.

11. The use of the expression of demethylase cDNA of claims 2 or 3 for the production of protein in vertebrate, insect or bacterial cells.

12. The use of claim 11 for producing antibodies against demethylase.

13. The use of the sequence of demethylase cDNA of claim 2 as a template to design antisense oligonucleotides and ribozymes.

14. The use of the predicted peptide sequence of demethylase cDNA of claim 2 to produce polyclonal or monoclonal antibodies against demethylase.

15. The use of expression of cDNA of claim 2 or 3 in two hybrid systems in yeast to identify proteins interacting with demethylase for diagnostic and therapeutic purposes.

16. The use of expression of cDNA of claim 2 or 3 in bacterial, vertebrate or insect cells to produce large amounts of demethylase for high throughput screening of

demethylase inhibitors for therapeutics and biotechnology and for obtaining the x-ray crystal structure.

17. A volatile assay for high throughput screening of demethylase inhibitors as therapeutics and anticancer agents which comprises the steps of:

- a) using transcribed and translated demethylase cDNA of claim 2 or 3 *in vitro* to convert methylcytosine present in methylated DNA samples to cytosine present in DNA and volatilize methyl group;
- b) determining the absence or minute amount of volatilized methyl group as an indication of an active demethylase inhibitor.


18. A volatile assay for the diagnostics of cancer in a patient sample which comprises the steps of:

- a) determining demethylase activity in patient samples by determining conversion of methylcytosine present in methylated DNA to cytosine present in DNA and volatilization of the methyl group released as methanol;
- b) determining the presence or minute amount of volatilized methyl group as an indication of cancer in said patient sample.

19. Use of an antagonist or inhibitor of DNA demethylase of claim 1 or 2 for the manufacture of a medication for cancer treatment, for restoring an aberrant methylation pattern in a patient DNA, or for changing a methylation pattern in a patient DNA.

20. Use according to claim 19, wherein said antagonist is a double stranded oligonucleotide that inhibits demethylase at a K_i of 50nM.

21. Use according to claim 20, wherein said oligonucleotide is



22. Use according to claim 19, wherein the inhibitor comprises an anti-DNA demethylase antibody or an antisense oligonucleotide of DNA demethylase or a small molecule.

23. Use according to one of claims 19 or 22, wherein the change of the methylation pattern activates a silent gene.

24. Use according to claim 23, wherein the activation of a silent gene permits the correction of genetic defect.

25. Use according to claim 24, wherein said genetic defect is β -thalassemia or sickle cell anemia.

26. Use of the demethylase of claim 1, for removing methyl groups on DNA in vitro.

27. Use of the demethylase of claim 1 or its cDNA of claim 2, for changing the state of differentiation of a cell to allow gene therapy, stem cell selection or cell cloning.





28. Use of the demethylase of claim 1 or its cDNA, of claim 2 for inhibiting methylation in cancer cells using vector mediated gene therapy.

29. An assay for the diagnostic of cancer in a patient, which comprises determining the level of expression of DNA demethylase of claim 1 in a sample





from said patient, wherein overexpression of said DNA demethylase is indicative of cancer cells.

003060 11112500


1/50



1A



SUBSTITUTE SHEET (DI 11 E 25)









FIG. 2C


FIG. 2D

5/50

005060-11115560


Fig. 3AFig. 3B




6/50




Labeled nucleotide:
[α 32P]-dCTP

Labeled nucleotide:
[α 32P]-dGTP






8/50




9/50




F I I I - 4 E

10/50



TSS-4F


11/50



$$\Delta G_{\text{Reaction}} = (93) - (103 + 91.5) = (-101.5) \text{ Kcal}$$


 11/50

12/50





TESS-BA


13/50




14/50



15/50



16/50



F18-70

F18-70

17/50

[illegible][illegible]

FILE - BA


MDCPALPCGW KKEEVIRKSG LSACKSDVYV FSPSGKKFRS 40
 KPOLARYLGN TVDLSSFDPR TGRMMPSKLG KNKQRLRNDP 80
 LLNQNGKGPDL NTTLPIRQTA SIFKQPVTKV TNHPSNKVKS 120
 DPQRMNEQPR QLFWEKRLQG LSASDVTEQI IKTMELPKGL 160

VVLNLTSQLC	KAFIVTDEDI	RKOERVOOV	RKILEDALMA	240
DILSRAADTE	EMDIEMDSGD	EA		262

711-1111

18/50

005060-11-11-5560



7-11-80

Human DNA demethylase cDNA-dMTase1 and predicted amino acid sequence

5'ggggcgctgg ccccgagaag gcgagacaa gatggccgcc catagcgctt ggaggaccta
 agaggcggtg gccggggcca cgcccgggc aggaggcccg ctctgtgcgc gcccgctcta
 tgatgcttgc gcgctcccc cgcgccgcg gctcggggcg gggcgggtct cccggattcc
 aagggtctcg ttacggaaga agcgcagcgc cggctgggga ggggctgga tgcgcgcga
 cccggggga gccgctgct gccggagca ggaggaggcg gagagtgcgg cgggcggcag
 cggcgtggc ggcgactccg ccatagagca ggggggccag ggcagcgcg tcgccccgtc
 cccggtgagc ggcgtgcga gggaaggcg tcggggcggc gccgtggcc gggggcggtg
 gaagcaggcg gccggggcg gcgctctg tggcgtggc cggggccggg gccgtggcg
 gggacggga cggggccgg gccggggcg cggcgtccc ccgagtggcg gcagcggcct
 tggcggcgac gccggcggt gccggcgcg cggcagcggc gccggcgcg cccccggcg
 ggagccggtc ccttcccgat aagagatgg attgcccgc cctcccccc ggatggaaag
 ggagagcggg aagagatgg attgcccgc cctcccccc ggatggaaag aggaggaaagt
 gatccgaaaa tctgggctaa gtgctggcaa ggcgatgtc tactacttca gtccaagtgg
 taagaagtgc agaagcaagc ctcagttggc aaggtacctg ggaaatactg ttgatctcag
 cagttttgac ttcagaactg gaaagatgat gcctagtaaa ttacagaaga acaaacagag
 actgcgaac gatccttca atcaaaataa gggtaaacca gacttgaata caacattgcc
 aattagacaa acagcatcaa tttcaaaaca accggttaacc aaagtcacaa atcatcctag

19/50

735-9A

20/50

taataaaagtg aaatcagacc cacaacgaat gaatgaacag ccacgtcagc ttttctggga
gaagaggcta caaggactta gtgcatcaga tgtaacagaa caaattataa aaaccatgga
actaccctaaa ggtcttcaag gagtgtgctc aggtagcaat gatgagaccc ttttatctgc
tgttgccagt gctttgcaca caagctctgc gccaatcaca gggcaagtct ccgctgctgt
ggaagaagaac cctgctgttt ggcttaacac atctcaaccc ctctgcaaa gctttattgt
cacagatgaa gacatcagga aacaggaaga gcgagtacag caagtacgca agaaattgga
agaagcactg atggcagaca tcttgtcgcg agctgctgat acagaagaga tggatatgga
aatggcacagt ggagatgaag cctaagaata tgatcaggta actttcgacc gactttcccc
aagrgaaaat tcctagaaat tgaacaaaaa tgtttccact ggcttttggc tgtaagaaaa
aaaatgtacc cgagcacata gagcttttta atagcactaa ccaatgcctt tttagatgta
tttttgatgt atatatctat tattcaaaaa atcatgttta ttttgagtcc taggacttaa
aattagtctt ttgtaatatc aagcaggacc ctaagatgaa gctgagcttt tgatgccagg
tgcaatctac tggaaatgta gcacttacgt aaaacatttg tttcccccac agttttaata
agaacagatc aggaattcta aataaatttc ccagttaaag attattgtga cttcacgtta
tataaacata tttttatact ttattgaaag gggacacctg tacattcttc catcatcact
gtaaaagacaa ataaatgatt atattcaca aaaaaaaaaa aaaaaa 3'

SEQ ID NO:1

21/50

MRAHPGGGCCPEEEGESAAGSGAGGDSAIEQGGQGSALAPSPVSGVR
 REGARGGGRGRWKQAGRGGVCGRGRGRGRGRGRGRGRGRGRPPSG
 GSGLGDDGGCGGGGGGAPRRREPVPFPSPGSAGPGPRGRPRATESGKRM
 DCPALPPGWKKEEVIRKSGLSAGKSDVYFSPSGKKFRSKPQLARYLGNT
 VDLSSDFRTGKMMPSKLQKNKQRLRNDPLNQNKGKPDNLTTLPPIRQTAS
 IFKQPVTKVTNHP SNKVKSDPQRMNEQPRQLFWEKRLQGLSASDVTEQII
 KTMELPKGLQGVGPGSNDETLLSAVASALHTSSAPITGQVSAAVEKNPAV
 WLNTSQPLCKAFIVTDEDIRKQEEERVQQVRKKLEALMADILSRAADTEE
 MDIEMDSGDEA

SEQ ID NO:2

009060-11115560

Human DNA demethylase homologue-dMTase2 and predicted amino acid sequence

5' agcgggcccga ggagcggggc gcaatggagc ggaagaggtg ggagtgcgcg gcgctcccg
 aggcctggga gaggaagaa gtgccagaa ggtcggggtc gtgcggcgcg cacagggatg
 tcttttacta tagcccgagc gggaagaagt tccgcagcaa gccgcagctg gcgcgctacc
 tgggcggctc catggacctg agcaccttcg acttcgcac gggcaagatg ctgatgagca
 agatgaacaa gagccgccag cgcgtgcgtc acgactcctc caaccaggtc aagggcaagc
 ccgacctgaa cacggcgctg cccgtgcgc agacggcgtc catcttcaag cagccgggtga
 ccaagattac caaccacccc agcaacaagg tcaagagcga cccgcagaag gcggltggacc
 agccgcgcaa gctcttctgg gagaagaagc tgagcggcct gaacgccttc gacattgctg
 aggaagtgtt caagaccatg gacctccca agggcctgca ggggggtggga cctggctgca
 cggatgagac gctgctgtcg gccatcgca gcgccctgca cactagcacc atgccccatca
 cgggacagct ctgcggccgc gtggagaaga acccggcgt atggctcaac accacgcagc
 ccctgtcaa agccttcctg gtgaccgacg aggacatcag gaagcaggaa gagctggctc
 agcaggtgcg gaagcggctg gaggagcgc tgatggccga catgctggcg cacgtggagg
 agctggcccc tgacggggag gcgccgtgg acaaggcctg cgctgaggac gacgacgagg
 aagacgagga ggaggaggag gaggagcccc acccggacc ggagatggag cacgtctagg
 gcagaggccc tgccgagagc ccgtgctgcc tgctggagcc gcctgcagac gcggtcctcg
 gcccacgtg aaccaggctc ggcggcggaag cccagccttg gagacacca ggaggaaaggc
 cgtgctcctg gctccctcct cggcccgctc ccaactcccg gggcctcggg gcacacagct
 ggggctgccc ccaccggaaa gacctccac gctcgtcctc tacagagtcc ggcttcggga
 agtgccgggt gctcctgggc cctgcctggc tccctacgac ctctgggctc gaggccagct

TTTT - 90

23/50

cctccccatg ccgctgtccc cagctccttg agactggaga gcagccagca ggtgcccgcc
 agctcggcgc cactgcttgc tgacagctgg gagggtttct cggctggag gcgtagtttt
 gaaactcaca tcaccactg tgcagcgtga ggaaggact ctggtctgct gtggggggca
 tgcaggacgg cgcaactctc tgccttgcca tgcgctgggt ggtgccacag agcctcaccg
 tgcctgagtg gcgtgcccag gaggccgct ctcttcagt aaatgtaaca cagtcgaggc
 acgtcatcgg gcagccttcc ctgttgcca acgccagcct tcgcttctga aaacaaaat
 ccagccgctg ccagtggga ctgtgtccc cggcgtgccc agaatgctcc actgccagcc
 ggccccctg cctcggttc cttctgttt agtggcgaca caggcaccca gctttgggg
 ggtgctgacg ctcccagggg tgcaggagc cactgggaca ggtgaggct ccagacgct
 cctcagagtg ccagctctc cagggagct ctggcccagg gcgttcttga gggatctgct
 ccttaacccc ccagtgcctt ggcgaggga ggttccaaag cacagagcc tggcccagt
 gacttttgcg gccagtccct ggtgcccctg ctgggcccctg ctggcccagt gagggttcc
 aacgggtggg ttcawtggcc tggcccavag gagccccac ctgcattgac cttagggcca
 tagagaggc ctgtcccgt gctgcccag ccaaggatct ggtcgtgcc ccagggggac
 tgatgggcaa gactgcccc tgtggctgga ctgtgacct gccctgagg gcctgaccgc
 gggagctgag gaagcgccg tccacctct gccctccag gaccgcattg gaggcagtgg
 gctggcagct tcctgtgct ccctgtcaga gtcaaaagcac aaatcctcag gacgggctca
 agggccaggg cagccgaggg aagctccagg tggggaccac gtcttcttga ggttggtgcc
 cactggctgg gacccttgc agtggggtgg cctccctct gtctgcctgg tggaggagac
 cgtgggctg gggacgtgac tgaataaagc caccatgggt ggatgtgctt gg^{3'}

SEQ ID NO:3

~~FILE~~ - DE

Lipman-Pearson Protein Alignment

Kluple: 2; Gap Penalty: 4; Gap Length Penalty: 12

Seq1(1>411)

Seq2(1>291)

human dMTase1 protein

human dMTase2 protein

Similarity

Gap

Gap

Consensus

(148>397)

(4>253)

76.0

0

0

250

(148>397)

(4>253)

76.0

0

0

250

v150 v160 v170 v180 v190 v200 v210

KRMDCPALPPGWKKEEVIRKSGISAGKSDVYFSPSGKKFRSKPQLARYLGNIVDLSSDFRIGKNMPSK
 KR:CPALP.GW.:EEV R:SGLSAG..DV:Y:SPSGKKFRSKPQLARYLG.:DLS:FDFRIGKM:SK
 KRWECPALPQGWEREVEVPRRSGLSAGHRUVFYFSPSGKKFRSKPQLARYLGSGMDLSTFDFRIGKMLMSK

25/50

^10

^20

^30

^40

^50

^60

^70

v220

v230

v240

v250

v260

v270

v280

LQKNKQRLRNDPLNQNKGKPDNLITLPIRQIASLIFKQFVTKVTNHPSNKKVKSDFQRMNEQPPQLFWKKRL
 :K::QR:R D: NQ KKKPDINT:LP:RQIASLIFKQFVTK:TNHPSNKKVKSDFQ:QPPQLFWKK:L
 MNKSRQFVRYDSSNQKGGKPDNIALFVRQIASLIFKQFVTKTNHPSNKKVKSDFQKAVDQPPQLFWKKL

^80

^90

^100

^110

^120

^130

^140

 77-96

26/50

v290 v300 v310 v320 v330 v340 v350
 QG LSASJVTIEQILIKIMELPKGLQG VGRG S NDEI LLSA VASALHTSSAPITIGQVSAAVEKNP A V W L N I T S Q P
 GL: A D::E::KIM: L P K G L Q G V G P G.. D E I L L S A: A S A L H T S: P I T G Q: S A A V E K N P: V W L N I T: Q P
 S G L N A F D I A F E L V K I M D L P K G L Q G V G R G C I D E I L L S A I A S A L H T S I M P I T G Q L S A A V E K N P G W L N I T Q P
 ^150 ^160 ^170 ^180 ^190 ^200 ^210

v360 v370 v380 v390
 L C K A F T V I D E D I R K Q E E R V Q Q V R K K L F F A I M A D I L S R A A D
 L C K A F: V I D E D I R K Q E E V Q Q V R K: L F F A I M A D: L: : : :
 L C K A F W I D E D I R K Q E E L V Q Q V R K R L F F A I M A D M L A H V E E
 ^220 ^230 ^240 ^250

F I S S E D
 - 9 H

Mouse DNA demethylase-dMTase1 and predicted amino acid sequence

5' ccgctctgcg ggcggggcgg gtctccggga ttccaaggcg tcggttacgg aagaagcgca
 gagccggctg gggagggggc tggatgcgc cgcacccggg gggagccgc tgcctgcccg
 agcaggagga ggggagagc gcggcgggcg gcagcgcg tcggcgcgac tccgccatag
 agcagggggg ccagggcagc gcgctcgctc cgtccccggt gagcgcgctg cgcagggaag
 gcgctcgggg cggcgccgct ggcggggggc ggtggaagca ggcggcccg ggcggcgggc
 tctgtggcgg tggccgtggc cgtggccggg gtccggggcg tcggccgggc cggggcccg
 gccggcgcg tcccagagt ggcggcagcg gccttggcgg cgacggcgcg ggcggcgcg
 gcggctgcgg cgtcggcagc ggtggcgcg tcgcccccg gcgggatacct gtcccttcc
 cgtcggggag ctcggggcg gggcccgagg gacccggcg cagcgagagc ggaagagga
 tggactgcc ggcctccc cccggatgga agaaggagga agtgatccga aaatcagggc
 tcagtctgg caagagcgat gtctactact tcagtcgaag tggtaagaag ttcagaagta
 aaactcagct ggcaagatc ctgggaaatg ctgttgacct tagcagtttt gacttcagga
 ccggcaagat gatgctagt aaattacaga agaacaagca gagactccg aatgaccccc
 tcaatcagaa caagggtaaa ccagacctga acacaacatt gccaattaga caaactgcac
 caattttcaa gcaaccagta accaaattca cgaaccacc gagcaataag gtgaagtacg
 accccagcg gatgaatgaa caaccagtc agcttttctg ggagaagagg ctacaaggac
 ttagcgcatc agatgtaaca gaacaattta taaaaacat ggagctacct aaaggtcttc
 aaggagtggg tccaggtagc aatgacgaga cccttctgtc tgcgtggtggc agtgccttac

115560 - 91

27/50

28/50

acacaagctc tgcgccatc acaggacaag tctctgctgc cgtgaaaag aacctgctg
 ttggccttaa cacatctcaa cccctctgca aagctttcat tgttacagat gaagacatta
 ggaacacagga agagcgagtc caacaagtac gcaagaaact ggaggaggca ctgatggcgg
 acatcctgtc ccgggctgcg gacacggagg aagtagacat tgacatggac agtggagatg
 aggcgtaaga atatgatcag gtaactttcg actgaccttc cccaagagca aattgctaga
 aacagaatta aaacatttcc actgggttcc gcctgtaaga aaaagtgtac ctgagcacat
 agctttttaa tagcactaac caatgccttt ttagatgtat ttttgatgta tatatctatt
 attccaaatg atgtttattt tgaatcctag gacttaaaat gagtctttta taatagcaag
 cagggccctt ccggtgcagt gcagcttlga gcccagggtgc agtctactgg aaaggtagca
 cttacgtgaa atatttgttt cccccacagt ttaatatata acagatcagg agtaccacaa
 aagtttccca attaaagatt attatacttc actgtatata acagattttt tatactttat
 tgaagaaga tacctgtaca ttcttccatc atcactgtaa agacaaaataa atgactatat
 tcac 3'

SEQ ID NO:5

FILE - 91

29/50

009060-11115560

MRAHPGGRCCEEGESAAGSGAGGDSAIEQGQGSALAPSPVSGVR
 REGARGGGRGRGWKQAAARGGVCGRGRGRGRGRGRGRGRGRPRQSG
 GSGLGDDGGGAGGCGVGSGGGVAPRRDPVPPSGSSGPGPRGPRATESG
 KRMDCPALPPGWKKEEVIRKSGLSAGKSDVYFSPSGKKFRSKPQLARYL
 GNAVDLSSFDFTGKMMPSKLQKNQRLRNDPLNQNKGPDLNLTTLPIRQ
 TASIFKQPVTKFTNHPSNKKVKSDDPQRMNEQPRQLFWEKRLQGLSASDVTE
 QIIKTWELPKGLQGVGPGSNDETLLSAVASALHTSSAPITGQVSAAVEKN
 PAVWLNTSQPLCKAFIVTDEDIRKQEERVQVRKKLEELMADILSRAAD
 TEEVDIDMSGDEA

SEQ ID NO:6



Mouse DNA demethylase-dMTase2 and predicted amino acid sequence

5' caccgcgcgg cgggtgggcg gagcggcccc cctagcgggg gctgtgaagc gcggggaggg
 ggccgagcgg gtggcgaagc cggcgcgcg cggctgggg gcggaggcgc gaggcccggtg
 ggacagaaca gctcggcga gtggcggcg cggagggagc cgaatcgcg acgagcccg
 gggtcgaac ttgcagaagc ggcgcgcg ggcgcatcg ccacggcgg ccgaaaagcc
 gggcgcaat ggagcggaag aggtgggagt gcccggcgct ccgcagggc tgggaaagg
 aagaagtgc caggagtcg gggctgtcg ccggccacag ggatgtctt tactatagcc
 ccagcgggaa gaagtccgc agcaagccac aactggcacg ttacctgggc ggatccatgg
 acctcagcac ctgcacttc cgcacggaa agatgttgat gaacaagatg aataagagtc
 gccagcgtgt gcgctatgat tcttccaacc aggtcaaggg caagcctgac ctgaacacc
 cgctgcctgt acggcagact gcattccat tcaagcaacc ggtgaccaag atcaccaacc
 accccagcaa caaggtcaag agcaccgc agaaggcagt ggaccagccg aggcagcttt
 tctgggagaa gaagtaagt ggattgagt cctttgacat tgcagaagaa ctggtcagga
 ccatggactt gcccaaggc ctgcagggag tgggccttg ctgtacagat gagcgtgc
 tgtcagccat tgcgagtgt ctacacaca gcacctgcc cattacaggc cagctctctg
 cagccgtgga gaagaacct ggtgtgtggc tgaacactgc acagccactg tgcaaacct
 tcatggtgac agatgacgac atcaggaaagc aggaggagct ggtacagcag gtacggaagc
 gcctggagga ggcactgat gccgacatgc tagtcatgt ggaggagctt gcccgagacg
 gggaggcacc actggacaag gcctgtgcag aggaggaa gaaggaggag

715560 - 91

30/50

31/50

aagagccgga gccagagcga gtgtagcaca ggtgcctgc ccaagtctgg gctgcagact
gccttcagcc ttgcctggac caggtagggg ccagacctgt aggaggcagc cgtccacctc
ctttccaaag cctcctgctt ccaggtctca gtgcaggagg cccctgtgga ccttgaactc
acttgtccct gcgctgcctg gcaggaaagg ccacactgaa agcagatgag cagtgaacca
actgagaggc cacctggaca cagtcacctc cctgcctcct tatcatagga caaggccttg
cttggcaccg aggagctggg agccgtgttg ggtgctggag gaagtttctg gaaacacacc
tggtatgcc caccttatgt ccctaaggct attacaggcc agggtttggg ctgctccggc
ccacagggct gccagcctc ccacactga ggttcagcag cccaccagga agtcacttcc
cttcaataaa ctgatggtag gaacttgtg 3'

SEQ ID NO:7



09/554414

32/50

MERKRWECPALPQWEREEVPRRSGLSAGHRDVFYVYSPSGKKFRSKPQLA
 RYLGGSMDLSTFDFTGKMLMNMKNKSRQVRVYDSSNQVKGKPDNLNTALP
 VRQTASIFKQPVTKITNHP SNKVKSDPQKAVDQPRQLFWEKKLSGLSAFD
 IAEELVRTMDLPKGLQGVPGCTDETLLSAIALHTSTLPI TGQLSAAV
 EKNPGVWLNTAQPLCKAFMVTDDDIRKQEELVQQVRKRLEEALMADMLAH
 VEELARDGEAPLDKACAEVEEEEEEEEEPEPERV

SEQ ID NO: 8



Lipman-Pearson Protein Alignment

Ktupe: 2; Gap Penalty: 4; Gap Length Penalty: 12

Seq1(1>414)	Seq2(1>285)	Similarity	Gap	Gap	Consensus
mouse dMTase1 protein	mouse dMTase2 protein	Index	Number	Length	Length
(151>400)	(4>253)	75.2	0	0	250
(151>400)	(4>253)	75.2	0	0	250
v160	v170	v180	v190	v200	v210
KRMDCPALPGWKKKEVIRKSCLSACKSDVYFSPSGKKFRSKPQLARVLGNVADLSSFDFTGKMMPSK					
KR : CPALP.GW.:EEV R:SGLSAG..DV:Y:SPSGKKFRSKPQLARVLG.:DLS:FDFTGKMM: :K					
KRWECPALPQGWEREVEVPRRSGLSAGHRDVFYFSPSGKKFRSKPQLARVLGGSMDLSTFDFTGKMMLNK					
^10	^20	^30	^40	^50	^60
					^70
v230	v240	v250	v260	v270	v280
LQKNKQRLRNDPFLNQNKGKPDINTILPIRQFASIFKQPVTKFTNHPSNKKVKSDFQRMNEQPRQLFWEKRL					
:K::QR:R D: NQ KGKPDINT:LP:RQIASIFKQPVTK:TNHPSNKKVKSDFQ: :QPRQLFWEK:L					
MNKSQRQVRVYDSSNQVKGKPDINTALFVRQIASIFKQPVTKFTNHPSNKKVKSDFQKAVDQPRQLFWEKKL					
^80	^90	^100	^110	^120	^130
					^140

33/50

v290


7355-90

34/50

	v370	v380	v390	v400
LCKAF:IVTDEDIPKQEEKVQQVRKKLEEFALMADII.SRAAD				
LCKAF:VTD:DIPKQEE VQQVRK: LEEALMAD:L:::				
LCKAF:VTDDIPKQEELVQQVRKRLEEFALMALMLAHVEE				
^220 ^230 ^240 ^250				

715 99

35/50



mistrane
dMTase







FIG. 10A

36/50


FIG. 10B

37/50


005060-11115560




39/50



40/50

FIG. 11F

41/50



— Origin

– Origin

FILE - 72A

000000 11115560







WO 99/24583

PCT/CA98/01059

09/554414

42/50

FILE-12B



43/50





FIG - 13A






FIG - 13B

44/50


7-11-13C

45/50


dMTase



18s





dMTase / 18s





TH-130



47/50


FIG. 14BFIG. 14C

48/50




- 15

49/50

Fig. 16

50/50

FIG. 17

**COMBINED DECLARATION FOR UTILITY OR DESIGN
PATENT APPLICATION (37 CFR 1.63) AND POWER OF ATTORNEY**

I hereby claim the benefit under 35 U.S.C. 120 of any United States application(s), or 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application.

U.S. Parent Application or PCT Parent Number	Parent Filing Date (MM/DD/YYYY)	Parent Patent Number (if applicable)
PCT/CA98/01059	11/12/1998	

☐ Additional U.S. or PCT International application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto:

As a named inventor, I hereby appoint the following registered practitioner(s) to prosecute this application and to transact all business in the Patent Trademark Office connected therewith:

☐ Customer Number
Place Customer Number Bar Code Label Here

OR ☒ Registered practitioner(s) name/registration number listed below

Name	Registration Number	Name	Registration Number
Joseph D. Evans	<u>26,269</u>	Herbert I. Cantor	<u>24,392</u>
Donald D. Evenson	<u>26,180</u>	Gary R. Edwards	<u>31,824</u>
Robert Mitchell	<u>25,007</u>	Guy Houle	<u>24,971</u>
Kevin P. Murphy	<u>26,674</u>	Robert Carrier	<u>30,726</u>
Michel J. Sofia	<u>37,017</u>	François Nadeau	<u>37,570</u>
Franco Côté	<u>37,037</u>		

Direct all correspondence to ☐ Customer Number OR ☒ Correspondence address below

Name Joseph D. Evans
 Address EVENSON, MCKEOWN, EDWARDS & LENAHAN
 Address Suite 700, 1200 G Street, N.W.
 City Washington State D.C. Postal Code 20005
 Country USA Telephone (202) 628-8800 Fax (202) 628-8844

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Name of sole or First Inventor:

☐ A petition has been filed for this unsigned inventor

Given Name (first and middle (if any))

Family Name or Surname

1-00 Moshe SZYF
 Inventor's Signature Moshe SZYF Date 28.6.2000
 Residence: City Côte St-Luc State Québec Country Canada Citizenship Canadian
 Post Office Address 7522 Wavell

City Côte St-Luc Province or State Québec Postal Code H4W 2L7 Country Canada

☒ Additional inventors are being named on the supplemental Additional Inventor(s) PTO/SB/02A attached hereto.

COMBINED DECLARATION FOR UTILITY OR DESIGN
PATENT APPLICATION (37 CFR 1.63) AND POWER OF ATTORNEY

PTO/SB/02A (3-97)

DECLARATION

ADDITIONAL INVENTOR(S)
Supplemental Sheet
Page 3 of 3

Name of Additional Joint Inventor, if any:

☐ A petition has been filed for this unsigned inventor

Given Name (first and middle (if any))

Family Name or Surname

2-00 Sanjoy BHATTACHARYA

Inventor's Signature [Signature] Date July 6/2000

Residence: City Montréal State Québec Country Canada Citizenship Indian CAX

Post Office Address Apartment 1403

2150 Mackay Street

City Montréal Province or State Québec Postal Code H3G 2M2 Country Canada

Name of Additional Joint Inventor, if any:

☐ A petition has been filed for this unsigned inventor

Given Name (first and middle (if any))

Family Name or Surname

3-00 Shyam RAMCHANDANI

Inventor's Signature [Signature] Date July 10/2000

Residence: City Niagara Falls State Ontario Country Canada Citizenship Canadian CAX

Post Office Address 5843 Hillcrest Crescent

City Niagara Falls Province or State Ontario Postal Code L2J 2A8 Country Canada

Name of Additional Joint Inventor, if any:

☐ A petition has been filed for this unsigned inventor

Given Name (first and middle (if any))

Family Name or Surname

Inventor's Signature _____ Date _____

Residence: City _____ State _____ Country _____ Citizenship _____

Post Office Address _____

City _____ Province or State _____ Postal Code _____ Country _____

☐ Additional inventors are being named on the supplemental Additional Inventor(s) PTO/SB/02A attached hereto.

SEQUENCE LISTING

<110> MCGILL UNIVERSITY
SZYF, Moshe
BHATTACHARYA, Sanjoy K.
RAMCHANDANI, Shyam

<120> DNA DEMETHYLASE, THERAPEUTIC AND
DIAGNOSTIC USES THEREOF

<130> 1770-183"PCT" FC/ld

<150> CA 2,220,805

<151> 1997-11-12

<150> CA 2,230,991

<151> 1998-05-11

<160> 10

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 1804

<212> DNA

<213> Unknown

<400> 1

ccgctctcgc	ggcggggcgg	gtctccggga	ttccaagggc	tcgggttacgg	aagaagcgca	60
gagccggctg	gggagggggc	tggatgcgcg	cgcaccggcg	gggaggccgc	tgctgcccgg	120
agcaggagga	gggggagagc	gcggcgggcg	gcagcggcgc	tggcgggcgc	tccggccatg	180
agcagggggg	ccaggggcgc	gcgctcgcct	cgtccccggc	gagcggcgctg	cgcagggaag	240
gcgctcgggg	cggcgggcgt	ggccgggggc	ggtggaaagca	ggcggcccg	ggcgcgcgcg	300
tctgtggccc	tggccgtggc	cgtggccggg	gtcggggcgc	tggccggggc	cggggccggg	360
gccgcggccc	tccccagagt	ggcggcagcg	gccttggcgg	cgacggcgcc	ggcgcgcgcg	420
gcggctgcgg	cgtcggcagc	ggtggcgcg	tcgccccccg	gcgggatcct	gtccctttcc	480
cgtcggggcg	ctcggggcgc	ggggccaggg	gacccccggc	cacggagagc	gggaagagga	540
tggactgccc	ggccctcccc	ccgggatgga	agaaggagga	agtgatccga	aaatcagggc	600
tcagtgcgtg	caagagcgat	gtctaactct	tcagtcgaag	tggtaagaag	ttcagaagta	660
aaacctcagct	ggcaagatgc	ctgggaaatg	ctgttgacct	tagcagtttt	gaacttcagga	720
ccggcgaagat	gatgcctagt	aaattacaga	agaacaagca	gagactccgg	aatgaccccc	780
tcaatcagaa	caagggtaaa	ccagacctga	acacaacatt	gccaataga	caaaactgcat	840
caattttcaa	gcaaacagta	acaaaattca	cgaaccaccc	gagcaataag	gtgaagtccg	900
acccccagcg	gatgaatgaa	caaccacgtc	agctttttctg	ggagaagagg	ctacaaggac	960
ttagcgcctc	agatgtaaca	gaacaaatta	taaaaacatt	ggagctacct	aaaggtcttc	1020
aaggagtcgg	tcagggtagc	aatgacgaga	cccttctgtc	tgctgtggcc	agtgctttac	1080
acacaagctc	tgcgccatc	acaggacaag	tctctgtctc	cgtggaaaaa	aacctgtctg	1140
tttggcttaa	cacatctcaa	ccctctcgca	aagctttcat	tgttacagat	gaagacatta	1200
ggaaaacagg	agagcagctc	caacaagtac	gcaagaaact	ggaggaggca	ctgatggccc	1260
acatctctgc	ccgggctgcg	gacacggagg	aagtagacat	tgacatagat	agtggaagatg	1320
aggcgtaaga	atatgatcag	gtaactttcg	actgaccttc	cccaagagca	aattgctaga	1380
aacagaaata	aaacattttc	actgggtttc	gcctgtaaga	aaaagtgtac	ctgagcacat	1440
agctttttta	tagcactaac	caatgccttt	tttagatgat	ttttgatgta	tatatctatt	1500
attccaaagt	atgtttattt	tgaatcctag	gacttaaaat	gagctcttta	taatatgcaag	1560
cagggccctt	ccgggtcagt	gcagctttga	ggccagggtc	agtcactcgg	aaaggtagca	1620
cttacgtgaa	atatttgttt	ccccacagt	tttaataata	acagatcagg	agtaccaa	1680

009000-ATTN:5260

aagtttccca attaaagatt attatacttc actgtatata aacagatttt tatactttat 1740
 tgaaagaaga tacctgtaca ttcttccatc atcactgtaa agacaaataa atgactatat 1800
 tcac 1804

<210> 2
 <211> 411
 <212> PRT
 <213> Unknown

<400> 2

Met Arg Ala His Pro Gly Gly Gly Arg Cys Cys Pro Glu Gln Glu Glu
 1 5 10 15
 Gly Glu Ser Ala Ala Gly Gly Ser Gly Ala Gly Gly Asp Ser Ala Ile
 20 25 30
 Glu Gln Gly Gly Gln Gly Ser Ala Leu Ala Pro Ser Pro Val Ser Gly
 35 40 45
 Val Arg Arg Glu Gly Ala Arg Gly Gly Gly Arg Gly Arg Gly Trp
 50 55 60
 Lys Gln Ala Gly Arg Gly Gly Gly Val Cys Gly Arg Gly Arg Gly Arg
 65 70 75 80
 Gly Arg Gly Arg Gly Arg Gly Arg Gly Arg Gly Arg Gly Arg
 85 90 95
 Pro Pro Ser Gly Gly Ser Gly Leu Gly Gly Asp Gly Gly Gly Cys Gly
 100 105 110
 Gly Gly Gly Ser Gly Gly Gly Ala Pro Arg Arg Glu Pro Val Pro
 115 120 125
 Phe Pro Ser Gly Ser Ala Gly Pro Gly Pro Arg Gly Pro Arg Ala Thr
 130 135 140
 Glu Ser Gly Lys Arg Met Asp Cys Pro Ala Leu Pro Pro Gly Trp Lys
 145 150 155 160
 Lys Glu Glu Val Ile Arg Lys Ser Gly Leu Ser Ala Gly Lys Ser Asp
 165 170 175
 Val Tyr Tyr Phe Ser Pro Ser Gly Lys Lys Phe Arg Ser Lys Pro Gln
 180 185 190
 Leu Ala Arg Tyr Leu Gly Asn Thr Val Asp Leu Ser Ser Phe Asp Phe
 195 200 205
 Arg Thr Gly Lys Met Met Pro Ser Lys Leu Gln Lys Asn Lys Gln Arg
 210 215 220
 Leu Arg Asn Asp Pro Leu Asn Gln Asn Lys Gly Lys Pro Asp Leu Asn
 225 230 235 240
 Thr Thr Leu Pro Ile Arg Gln Thr Ala Ser Ile Phe Lys Gln Pro Val
 245 250 255
 Thr Lys Val Thr Asn His Pro Ser Asn Lys Val Lys Ser Asp Pro Gln
 260 265 270
 Arg Met Asn Glu Gln Pro Arg Gln Leu Phe Trp Glu Lys Arg Leu Gln
 275 280 285
 Gly Leu Ser Ala Ser Asp Val Thr Glu Gln Ile Ile Lys Thr Met Glu
 290 295 300
 Leu Pro Lys Gly Leu Gln Gly Val Gly Pro Gly Ser Asn Asp Glu Thr
 305 310 315 320
 Leu Leu Ser Ala Val Ala Ser Ala Leu His Thr Ser Ser Ala Pro Ile
 325 330 335
 Thr Gly Gln Val Ser Ala Ala Val Glu Lys Asn Pro Ala Val Trp Leu
 340 345 350
 Asn Thr Ser Gln Pro Leu Cys Lys Ala Phe Ile Val Thr Asp Glu Asp
 355 360 365

009500 11115500

Ile Arg Lys Gln Glu Glu Arg Val Gln Gln Val Arg Lys Lys Leu Glu
 370 375 380
 Glu Ala Leu Met Ala Asp Ile Leu Ser Arg Ala Ala Asp Thr Glu Glu
 385 390 395 400
 Met Asp Ile Glu Met Asp Ser Gly Asp Glu Ala
 405 410

<210> 3
 <211> 1589
 <212> DNA
 <213> Unknown

<400> 3

cacgcgcggg cgggtggggg gagcgccccc cctagcgggg gctgtgaagc gcgggggagg 60
 ggccgagcgg gtggcggaag cggcgcgcg cgggctgggg gcgaggggcg gaggcccggt 120
 ggacagaaac gctgcggcga gtggcgcgcg cggaggagcg cgaatcgccg acgagcccg 180
 gggtcgcaac ttgcagaagc ggccgcgcg cggcgatcgg ccgcggcggg cggaaaagcc 240
 ggggcgcaat ggagcggaag aggtgggagt gcccggcgct ccgcgaggcg tggaaaagg 300
 aagaagtgcc caggaggtcg ggggtgtcgg ccggccacag ggaatgtctt tactatagcc 360
 ccagcgggaa gaagtccgc agcaagccac aactggcacg ttacctgggc ggatccatgg 420
 accctcagac cctcgacttc cgcaccggaa agatgttgat gaacaagatg aataagagtc 480
 gccagcgtgt gcgctatgat tcttccaacc aggtcaaggg caagcctgac ctgaacaccg 540
 cgctgcctgt acggcagact gcattccatc tcaagcaacc ggtgaccaag atcaccaccc 600
 accccagcaa caaggtcaag agcgacccgc agaaggcagt ggaccagccg aggcagcttt 660
 tctgggagaa gaagctaaat ggattgagt cctttgacat tgcagaagaa ctggtcagga 720
 ccatgtactt gcccaaggcg ctgcaggagg tgggcctcgg ctgtacagat gagacgtctg 780
 tgtcagccat tgcagtgct ctacacacca gcaccctgcc cattacaggc cagctctctg 840
 cagccgtgga gaagaacct ggtgtgtggc tgaacactgc acagccactg tcaaaagcct 900
 tcatgtgtgc agatgacgag atcaggaagc aggaggagct ggtacagcag gtacgggaagc 960
 gcctggagga ggcactgat gccgacatgc tagctcatgt ggaggagcct gcccgagacg 1020
 gggaggccacc actggacaag gcctgtgcag aggaggaaga ggaggaggaa gaggaggagg 1080
 aagagccgga gccagagcga gtgtagcaca ggtgcctcgc ccaagtctgg gctgcagact 1140
 gcccttcagc ttgcctggac caggtagggg ccagacctgt aggaggcagc cgtccacctc 1200
 ctttccaaag cctcctgctt ccaggtctca gtgcaggagg cccctgtgga ccttgaactc 1260
 acttgtccct gcgctgcctg gcagggaagc ccacactgaa agcagatgag cagtgaacca 1320
 actgagagcg cacctggaca cagtcacctc cctgcctcct tatcatagga caaggccttg 1380
 cttggaccgc aggagctggg agccgtgttg ggtgctggag gaagtcttcg gaaacacacc 1440
 tggctatgcc caccttatgt ccctaaggct attacaggcc agggtttgag ctgcctcggc 1500
 ccacagggct gccacgctc ccacactga gggctcagcag cccaccagga agtcactttc 1560
 cttcaataaa ctgatggtag gaacttgtg 1589

<210> 4
 <211> 291
 <212> PRT
 <213> Unknown

<400> 4

Met Glu Arg Lys Arg Trp Glu Cys Pro Ala Leu Pro Gln Gly Trp Glu
 1 5 10 15
 Arg Glu Glu Val Pro Arg Arg Ser Gly Leu Ser Ala Gly His Arg Asp
 20 25 30
 Val Phe Tyr Trp Ser Pro Ser Gly Lys Lys Phe Arg Ser Lys Pro Gln
 35 40 45
 Leu Ala Arg Tyr Leu Gly Gly Ser Met Asp Leu Ser Thr Phe Asp Phe
 50 55 60
 Arg Thr Gly Lys Met Leu Met Ser Lys Met Asn Lys Ser Arg Gln Arg
 65 70 75 80

0055444 000600

Val Arg Tyr Asp Ser Ser Asn Gln Val Lys Gly Lys Pro Asp Leu Asn
 85 90 95
 Thr Ala Leu Pro Val Arg Gln Thr Ala Ser Ile Phe Lys Gln Pro Val
 100 105 110
 Thr Lys Ile Thr Asn His Pro Ser Asn Lys Val Lys Ser Asp Pro Gln
 115 120 125
 Lys Ala Val Asp Gln Pro Arg Gln Leu Phe Trp Glu Lys Lys Leu Ser
 130 135 140
 Gly Leu Asn Ala Phe Asp Ile Ala Glu Glu Leu Val Lys Thr Met Asp
 145 150 155 160
 Leu Pro Lys Gly Leu Gln Gly Val Gly Pro Gly Cys Thr Asp Glu Thr
 165 170 175
 Leu Leu Ser Ala Ile Ala Ser Ala Leu His Thr Ser Thr Met Pro Ile
 180 185 190
 Thr Gly Gln Leu Ser Ala Ala Val Glu Lys Asn Pro Gly Val Trp Leu
 195 200 205
 Asn Thr Thr Gln Pro Leu Cys Lys Ala Phe Met Val Thr Asp Glu Asp
 210 215 220
 Ile Arg Lys Gln Glu Glu Leu Val Gln Gln Val Arg Lys Arg Leu Glu
 225 230 235 240
 Glu Ala Leu Met Ala Asp Met Leu Ala His Val Glu Glu Leu Ala Arg
 245 250 255
 Asp Gly Glu Ala Pro Leu Asp Lys Ala Cys Ala Glu Asp Asp Asp Glu
 260 265 270
 Glu Asp Glu Glu Glu Glu Glu Glu Pro Asp Pro Asp Pro Glu Met
 275 280 285
 Glu His Val
 290

<210> 5
 <211> 1966
 <212> DNA
 <213> Unknown

<400> 5
 gggggcggtg ccccgagaag gccgagacaa gatggcgcgc catagcgctt ggaggaccta 60
 agaggcggtg gccggggcca cggcccgggc agggaggcgc ctctgtgcgc gcccgctcta 120
 tgatgcttgc ggcggtcccc cgccgcgcgc gctgcggggc gggcggggtct ccgggattcc 180
 aaggcgctcg ttacggaaag agcgagcgcc cggttgggga gggggcgctgga tgcgcgcgca 240
 cccgggggga ggcgcgtgct gcccgagaca ggaggagggg gagagtgccg cggcgccgag 300
 cggcgctggc ggcgactccg ccatagagca gggggggcca ggcagcgccg tcgccccgtc 360
 cccggtgagc ggcgtgcgca gggaagggcg tcggggcgcc ggcggtggcc gggggcggtg 420
 gaagcaggcg gcccgggggc gcgcgctctg tggccgttgc cggggccggg gccgtggccg 480
 gggaggggga cggggccggg gccggggcgc cggcgctccc ccgagtgccg gcagcggcct 540
 tggcgcgac gccggcggtc gcggcgcgcc cggcagcggt ggcgagcgcc cccccggcg 600
 ggagcgggtc cctttccgct cggggagcgc gggggccggg cccaggggag cccggggcac 660
 ggagagcggg aagaggtatg attgcccggc cctccccccc ggatggaaga aggaggaagt 720
 gatccgaaaa tctgggctaa gtgctggcaa gaggcgatgc tactactta gtccaagtgg 780
 taagaagtgc agaagcaagc ctcagttggc aaggtacctg ggaataactg ttgatctcag 840
 cagttttgac ttcagaactg gaaagatgat gcctagttaa ttacagaaga acaaacagag 900
 actcggaacg gatcctctca atcaaaataa gggtaaacca gacttgaata caacattgcc 960
 aattagacaa acagcatcaa ttttcaacaa accggtaacc aaagtacaaa atcatcctag 1020
 taataaagt aaatcagacc cacaacgaat gaatgaacag ccacgtcagc ttttctggga 1080
 gaagagggcta caaggactta gtgcatacaga caaattataa aaacattgga 1140
 actaccacaa ggtcttcaag gattgtgtcc aggtagcaat gatgagacc ttttatctgc 1200
 tgttgccagt cctttgcaca caagctctgc gccatacaca gggcaagtct ccgctgctgt 1260
 ggaaaaaac cctgtctgtt ggcttaacac atctcaaccc ctctgcaaaag cttttattgt 1320

Thr Met Glu Leu Pro Lys Gly Leu Gln Gly Val Gly Pro Gly Ser Asn
 305 310 315 320
 Asp Glu Thr Leu Leu Ser Ala Val Ala Ser Ala Leu His Thr Ser Ser
 325 330 335
 Ala Pro Ile Thr Gly Gln Val Ser Ala Ala Val Glu Lys Asn Pro Ala
 340 345 350
 Val Trp Leu Asn Thr Ser Gln Pro Leu Cys Lys Ala Phe Ile Val Thr
 355 360 365
 Asp Glu Asp Ile Arg Lys Gln Glu Arg Val Gln Gln Val Arg Lys
 370 375 380
 Lys Leu Glu Glu Ala Leu Met Ala Asp Ile Leu Ser Arg Ala Ala Asp
 385 390 395 400
 Thr Glu Glu Val Asp Ile Asp Met Asp Ser Gly Asp Glu Ala
 405 410

<210> 7
 <211> 2392
 <212> DNA
 <213> Unknown

<400> 7

agcgggcccga ggagccgggc gcaatgggagc ggaagaggtg ggagtgcctcg gcgctcccgc 60
 agggctgggga gagggaagaa gtgcccagaa ggtcggggct gtccggccggc cacagggatg 120
 tcttttacta tagcccagagc gggaagaagt tccgcagcaa gccgcagctg cgcgcgtacc 180
 tgggcggctc catggacctg agcaccctcg acttccgcac gggcaagatg ctgatgagca 240
 agatgaacaa gagccggcag cgcgtgcgct acgactcttc caaccagatc aagggcgaagc 300
 ccgacctgaa cacggcgctg cccgtgcgcc agacggcgct catcttcaag cagccggtga 360
 ccaagattac caaccacccc agcaacaagg tcaagagcga cccgcagaag cgggtggacc 420
 agcccgcgcca gctctcttgg gagaagaagc tgacggcgct gaacgccttc gacattgctg 480
 agggagctggt caagaccatg gacctcccca agggcctgca ggggggtggga ctgggtcgca 540
 cggatgagac gctgctgtcg gccatcgcca gcgccttgca cactagcacc atgcccacatc 600
 cgggacagct ctgcggccgc gtggagaaga acccggcgct atggctcaac accacgcagc 660
 ccctgtgcaa agccttcatg gtgaccgacg aggcacatcag gaagcaggaa gagctgggtg 720
 agcaggtgctg gaacggctg gagggaggcg tgatggccga catgctggcg cacgtggagg 780
 agctggcccc tgacggggag gcgcccgtg acaaggcctg cgtgaggac gagcagcagg 840
 aagacagga gaggaggag gagggagccc acccgacccc ggagatggag cacgtctagg 900
 cgagaggccc tgccgagagc cctgtgctgc tgcgtggacc gcctgcagac cgggtctctg 960
 gccccacgtg aaccaggctc ggccgcgaag cccagccttg gagacaccaa ggaggaagcg 1020
 cgtgctctg gctccctctc cggcccgctc ccacttcccg gggcctcggg gcacacagct 1080
 ggggctgccc ccaccgaaa gacctccac gctcgtctc tacagagtcc ggctctggga 1140
 agtgcgggtg gctctcgggc cctgcctggc tccctacgac ctttgggtc gagggcagct 1200
 cctcccatct cccgctgtcc cagctccttg agactggga gcagccagca ggtgcccggc 1260
 agctcggcgc acgggttgc tgacagctgg gaggtttctt cggctggag cgtagtttt 1320
 gaaactcaca tcaccactg tgcagcgtga ggacgggact ctggctgct gtggggggga 1380
 tgcaggacgc gcacactct tgccctgcca tgcggctggt ggtgccacag agctccaccg 1440
 tgcttgagt gcgtgccac gagggccgct ctccctcagt aaatgtaaca cagtgcaggc 1500
 acgtcatcgc gcagccttcc ctgtgtgcca acgccaagct tcgctcttga aaaccaaat 1560
 ccagccgctg ccagtcggga cttgtgcgcc cggcgctgcc agaattgctc actgcacagc 1620
 gccccctct cctcgtttc cctctgttt agtggcgaca caggcaccca gctttggggt 1680
 ggtgctgaag ctcccagggg tgcccaggagc cactgggaca ggttgaggct ccacagcgtc 1740
 cctcgagggt cccagctctc caggagcctt ctggcccaag cgtttctga gggatctgct 1800
 ccttaacccc ccagctcctt ggccgaggga ggttccaa gcacagacgc tgccccagct 1860
 ggactttggc gccagtcctt ggggtgcttc ctggggccctg cttgcccagt gagggttctc 1920
 aacgggtggg ttcawtggcc tggcccavag ctagcccccac ctgcattgac cttaggcccca 1980
 tagagagggg ctgtcccggt gctgcgccag ccaaggtatc ggtcgtgcc ccagggggac 2040
 tgatgggcaa gactgcggcc tgtggctgga ctgtgaccat cctgatggg gcttgaccgc 2100
 gggagctgag gaagcgccgc tccacgtct gccctccaa gaccgcgatg gaggcagtg 2160

gctggcagct tcctgctgct cctgtcaga gtcaaagcac aaatcctcag gacgggctca 2220
 agggccaggg cagccgaggg aagctccagg tggggaccac gtcttcctga gggtgggtgcc 2280
 cactggctgg gaccctttgc agtggggtgg cctccctct gtctgcctgg tggagggagc 2340
 cgtgggctgg gggacgtgac tgaataaagc caccatgggt ggatgtgctt gg 2392

<210> 8

<211> 285

<212> PRT

<213> Unknown

<400> 8

Met Glu Arg Lys Arg Trp Glu Cys Pro Ala Leu Pro Gln Gly Trp Glu
 1 5 10 15
 Arg Glu Glu Val Pro Arg Arg Ser Gly Leu Ser Ala Gly His Arg Asp
 20 25 30
 Val Phe Tyr Tyr Ser Pro Ser Gly Lys Lys Phe Arg Ser Lys Pro Gln
 35 40 45
 Leu Ala Arg Tyr Leu Gly Gly Ser Met Asp Leu Ser Thr Phe Asp Phe
 50 55 60
 Arg Thr Gly Lys Met Leu Met Asn Lys Met Asn Lys Ser Arg Gln Arg
 65 70 75 80
 Val Arg Tyr Asp Ser Ser Asn Gln Val Lys Gly Lys Pro Asp Leu Asn
 85 90 95
 Thr Ala Leu Pro Val Arg Gln Thr Ala Ser Ile Phe Lys Gln Pro Val
 100 105 110
 Thr Lys Ile Thr Asn His Pro Ser Asn Lys Val Lys Ser Asp Pro Gln
 115 120 125
 Lys Ala Val Asp Gln Pro Arg Gln Leu Phe Trp Glu Lys Lys Leu Ser
 130 135 140
 Gly Leu Ser Ala Phe Asp Ile Ala Glu Glu Leu Val Arg Thr Met Asp
 145 150 155 160
 Leu Pro Lys Gly Leu Gln Gly Val Gly Pro Gly Cys Thr Asp Glu Thr
 165 170 175
 Leu Leu Ser Ala Ile Ala Ser Ala Leu His Thr Ser Thr Leu Pro Ile
 180 185 190
 Thr Gly Gln Leu Ser Ala Ala Val Glu Lys Asn Pro Gly Val Trp Leu
 195 200 205
 Asn Thr Ala Gln Pro Leu Cys Lys Ala Phe Met Val Thr Asp Asp Asp
 210 215 220
 Ile Arg Lys Gln Glu Glu Leu Val Gln Gln Val Arg Lys Arg Leu Glu
 225 230 235 240
 Glu Ala Leu Met Ala Asp Met Leu Ala His Val Glu Glu Leu Ala Arg
 245 250 255
 Asp Gly Glu Ala Pro Leu Asp Lys Ala Cys Ala Glu Glu Glu Glu Glu
 260 265 270
 Glu Glu Glu Glu Glu Glu Pro Glu Pro Glu Arg Val
 275 280 285

<210> 9

<211> 17

<212> DNA


<213> Unknown

<400> 9

ctggcaagag cgatgtc

United States Patent & Trademark Office

Office of Initial Patent Examination -- Scanning Division



Application deficiencies were found during scanning:

☒ Page(s) 10 of 3 of Declaration were not present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ Scanned copy is best available.